



DEMS WORKING PAPER SERIES

Strategy-proof preference aggregation and the anonymity-neutrality tradeoff

**Stergios Athanasoglou, Somouaoga
Bonkounou, Lars Ehlers**

No. 519 – April 2023 – Last revised April 2025

**Department of Economics, Management and Statistics
University of Milano – Bicocca
Piazza Ateneo Nuovo 1 – 2016 Milan, Italy
<http://dems.unimib.it/>**

Strategy-proof preference aggregation and the anonymity-neutrality tradeoff*

Stergios Athanasoglou[†] Somouaoga Bonkounou[‡] Lars Ehlers[§]

Abstract

Consider a setting in which individual strict preferences need to be aggregated into a social strict preference relation. For two alternatives and an odd number of agents, it follows from May's Theorem that the majority aggregation rule is the only one satisfying anonymity, neutrality and strategy-proofness (SP). For more than two alternatives, anonymity and neutrality are incompatible for many problem instances and we explore this tradeoff for strategy-proof rules. The notion of SP that we employ is Kemeny-SP (K-SP), which is based on the Kemeny distance between social orderings and strengthens previously used concepts in an intuitive manner. Dropping anonymity and keeping neutrality, we identify and analyze the first known nontrivial family of K-SP rules, namely semi-dictator rules. For two agents, semi-dictator rules are characterized by strong unanimity, neutrality and K-SP. For an arbitrary number of agents, we generalize semi-dictator rules to allow for committees and show that they retain their desirable properties. Dropping neutrality and keeping anonymity, we establish possibility results for three alternatives. We provide a computer-aided solution to the existence of a strongly unanimous, anonymous and K-SP rule for two agents and four alternatives. Finally, we show that there is no K-SP and anonymous rule which always chooses one of the agents' preferences.

Keywords: Preference aggregation, strategy-proofness, anonymity, neutrality, Kemeny

*We are grateful to two anonymous referees and the associate editor for useful comments and suggestions. The second author acknowledges financial support from the Swiss National Foundation, project number 100018 - 207722. The third author acknowledges financial support from SSHRC (Canada) under Insight Grant 435-2023-0129.

[†]Department of Economics, University of Milan - Bicocca, stergios.athanasoglou@unimib.it.

[‡]Department of Economics, University of Lausanne, bksom@gmail.com.

[§]Département de sciences économiques and CIREQ, Université de Montréal, C.P. 6128, Succursale Centre-Ville, Montréal, Québec H3C 3J7, Canada; e-mail: lars.ehlers@umontreal.ca.

1 Introduction

We study the problem of aggregating individual preferences into one preference representing society. Applications include elections where voters with individual preferences choose their political representatives, instances in which groups of agents (such as committees, governing bodies or consortia) must reach consensus on their collective preferences, and macroeconomic models that analyze the choices of a representative agent of the society. In all of the above situations the collective preference must possess the same properties as individual preferences and agents would like it to be “as similar as possible” to their own preference.

At an abstract level, the general preference-aggregation model assumes a set of agents and a finite set of alternatives A . Agents have (strict) preferences over A , modeled as binary relations on A . The exact form these preferences take varies and depends on the nature of the alternative set A . Once the individual preferences have been defined, a function transforms them into yet another preference relation on A , interpreted as society’s preference. This social preference is subsequently used to reach a collective decision. Examples of this process include sports or academic competitions in which judges render their individual judgments of the participants (expressed in ordinal or cardinal terms) which are then synthesized to obtain a collective judgment. This judgment may take various forms: the sole identity of the winner, a full ordering of the contestants, a partial ordering of the top performers, a partition of acceptable/unacceptable candidates, etc. Along related but also different lines, the output of the preference-aggregation process may provide a solution to the problem of collective choices when there is uncertainty on the available alternatives. For instance, in the academic job market, departments often need to fill an open position without knowing whether any given applicant will accept their offer. In this framework, an aggregate candidate ordering serves as a tool for selecting the most desirable candidate among those who are available.

Naturally, agents want the aggregate preference to be as similar as possible to their individual preferences. This consideration introduces a novel challenge: given two binary relations on A , how do we measure their degree of similarity? The answer to this question is not obvious. However, if we want to study preference aggregation from a mechanism-design perspective, we need to address this issue. Indeed, it is impossible to formulate a notion of non-manipulability for an aggregation rule unless we have a way of determining the closeness of a given binary relation on A to an agent’s own preferences. Such a measure of closeness acts as a proxy for desirability.

To deal with this issue, our paper adopts the Kemeny metric [30] as method of measuring the distance between two linear orderings of the alternatives¹. The Kemeny distance is grounded on a natural axiomatic foundation by Can and Storcken [21] and enables an intuitive strengthening of previous non-manipulability concepts in preference aggregation by Bossert and Sprumont [17] and Athanasoglou [6]. Kemeny-based preferences are an example of metrizable preferences over linear orders as discussed by Meskanen and Nurmi [38] and Laffond et al. [35]. In this setting, orderings are ranked via a distance function that satisfies metric conditions such that the desirability of an ordering for a given agent is decreasing in its distance from that agent’s own ordering of the alternatives.

Kemeny-based preferences hold economic and axiomatic appeal and represent a reasonable way of modeling preferences over orderings. At the same time, Nishimura and Ok [41] demonstrate that these benefits may come at the cost of questionable implications for choice behavior.² The interested reader should be mindful of this fact. An alternative approach would involve (i) imposing less detailed structure on agent preferences and (ii) allowing for greater heterogeneity where preferences over orders preserve the betweenness relation as a minimal consistency requirement.

The Kemeny distance induces single-peaked preferences as studied by Moulin [39], Demange [23], and Barberà et al. [12], among others. That is, each agent has a unique maximal order, his true preference relation, and orders that are more distant from this ideal in the Kemeny sense are less preferred. It is tempting to study strategy-proof and neutral rules on a broader single-peaked preference domain where the true preference relation remains the most

¹The Kemeny distance between two binary relations is defined as the number of pairs of alternatives at which they differ. For a precise definition see Section 3.3.

²This point is best illustrated with an example adapted by Nishimura and Ok [41]. Suppose there are three job market candidates, A, B and C, and Prof. Smith ranks A first, B second and C third (his ordering is denoted ABC). Suppose, further, that the departmental aggregate ordering of the candidates will be used to select the winning candidate among those that will actually turn out to be available. Reflecting complete ignorance on this score, the set of available candidates is assumed to be any member of the power set of $\{A, B, C\}$. Given a realization of the set of available candidates, the department picks the highest-ranked candidate within that set according to the departmental ordering (the so-called “maximal” element of the set). Now suppose Prof. Smith is confronted with two different candidate orderings that the department is contemplating: BAC and ACB. Which does he prefer? According to the Kemeny-based approach, Prof. Smith is indifferent between the two orderings as they both differ from his own preferences with respect to a single candidate pair. However, this conclusion might no longer hold if we take into account the possible realizations of the set of available candidates. For example, if the available candidates are all three A, B, and C, then when using BAC the department picks B while using ACB leads to A. Here, Prof. Smith would have preferred ordering ACB to BAC. A similar conclusion applies if the available candidates had been A and B. Conversely, Prof. Smith would have preferred ordering BAC to ACB if only candidates B and C had been available. In all other candidate availability scenarios, the two orderings yield identical recommendations and so the professor is indifferent between them. Tallying all of this information, we see that there are strictly more instances in which the professor prefers ACB to BAC than vice versa. Thus, it is reasonable to expect that, if given the choice, he will prefer the departmental ordering ACB to BAC.

preferred one, and where agents are allowed to have any preference relation over orders away from this ideal. This domain is richer than the domain induced by the Kemeny distance but would most likely result in an impossibility similar to Gibbard [24] and Satterthwaite [43]. For example, Bonkougou [16, Theorem 1] studied strategy-proof and neutral rules on the betweenness-consistent single-peaked domain and showed that allowing for any heterogeneity results in dictatorship. The Kemeny distance is arguably an important restriction on agent preferences, but it allows the construction of non-dictatorial strategy-proof rules.

Our contribution. Working within the preference-aggregation paradigm, our paper studies methods of selecting a collective (or social) ordering of the alternatives when individual preferences over them are themselves orderings.³ We refer to such procedures as *aggregation rules* or simply *rules*. This special case of the preference-aggregation framework, known as *Arrovian preference aggregation*, is a classic social-choice model pioneered by Arrow [5] in the early nineteen-sixties.

There are three canonical properties that aggregation rules aspire to satisfy: fairness, efficiency, and incentive-compatibility or, as is more commonly referred to in this literature, *strategy-proofness*. The fairness notions of anonymity and neutrality are intuitively appealing and enjoy widespread application. A rule satisfies *anonymity* if it is invariant to reshuffling agents’ identities, and thus treats agents equally when determining the collective outcome. By contrast, a rule satisfies *neutrality* if it does not systematically favor one alternative over another, which is a salient concern when the alternatives represent morally relevant entities.

In the job market analogy, anonymity would require that all faculty members have an equal say in the hiring process, regardless of seniority and area of expertise, whereas neutrality would translate to the requirement that candidates’ identities (and, by extension, their demographic, socioeconomic and cultural characteristics) not bias the rule’s outcome. Meanwhile, strategy-proofness would ensure that no faculty member has an incentive to strategically misrepresent their true ranking of the candidates, in the hopes of obtaining a departmental ordering that more closely aligns with their preferences.

When there are just two alternatives and an odd number of agents, May’s Theorem [37] establishes that majority rule is the only rule satisfying anonymity, neutrality and strategy-proofness.⁴ However, when there are more than two alternatives we show that, for many problem instances, there exists no rule that can simultaneously satisfy both anonymity and neutrality. This result is closely related to a similar finding by Moulin [40] regarding choice rules (which only choose a winning alternative and not an entire ordering). Evidently,

³The term “ordering” denotes a complete, reflexive, antisymmetric, and transitive binary relation.

⁴For a general number of agents, May’s characterization encompasses for multi-valued rules and substitutes strategy-proofness with an intuitive monotonicity requirement known as positive responsiveness.

obtaining May-like possibility results for strategy-proofness involves dropping either anonymity or neutrality. This is the path we pursue in the current paper.

The efficiency standard we impose is *strong unanimity* (where society’s preference shall respect any unanimous preference over two alternatives), an analogue of weak Pareto due to Arrow [5] that has been extensively used in the recent literature on preference aggregation. As far as non-manipulability is concerned, we adopt Kemeny strategy-proofness (K-SP) by using the intuitive and axiomatically-founded notion of Kemeny distance. This significantly strengthens previously used incomplete concepts based on the betweenness relation [17, 27, 6, 7]. Prior to our work there was no known nontrivial rule satisfying K-SP for all problem instances.

When we dispense with anonymity but keep neutrality, we propose and analyze the first known nontrivial (i.e., non-dictatorial and non-constant) family of K-SP rules, *semi-dictator* rules. As their name suggests, these rules grant outside influence to a single agent but stop short of being full dictatorships. They do so by incorporating voting by committees (Barberà et al. [14]) in a way that meaningfully restricts the power of the semi-dictator without violating K-SP. When there are just two agents, semi-dictator rules are characterized by strong unanimity, neutrality and Kemeny strategy-proofness. For an arbitrary number of agents, we show that semi-dictator rules retain their desirable properties.

When we dispense with neutrality but keep anonymity, we establish possibility results for the case of three alternatives. In particular, we show that certain subfamilies of Condorcet-Kemeny [17] and fixed-benchmark [7] rules satisfy anonymity, strong unanimity and K-SP. These results do not carry over when the number of alternatives exceeds three. To explore the case of four alternatives and two agents, we frame the rule-existence problem as an integer program. Consequently, we are able to identify computationally a strongly unanimous, anonymous and K-SP rule. This rule is, in some sense, similar to semi-dictator rules in that it grants special status to a unique “losing alternative”. This alternative is always placed as low as possible in the social ordering, subject to respecting strong unanimity, and the rule treats all other alternatives in a balanced (though non-neutral) fashion. Finally, we show that, for many instances, there is no anonymous and K-SP rule which always selects one of the agents’ preferences. This preference selection property is related to the requirement of “peak selection” often used in problems with one public good (Moulin [39] and others). We suspect that stronger impossibility results involving anonymity, K-SP and various notions of efficiency are likely to hold.

Related Work. Our paper is relevant for two strands of the broader social-theoretic literature. The first regards May’s Theorem and its various extensions when the number of

alternatives exceeds two, whereas the second deals with the nontrivial issue of how to model strategy-proofness in Arrovian aggregation.

We begin with the relevant literature on May’s Theorem in multi-alternative environments. We note that, unlike our paper, all the references we discuss deal with *choice* rules, as opposed to *aggregation* rules. Choice rules select a winning alternative, not an ordering of alternatives. Goodin and List [26] allowed for multi-valued rules and extended May’s result to the setting in which agents cast single-alternative votes among a set of more than two alternatives. In particular, they showed that majority rule is characterized by anonymity, neutrality and positive responsiveness in this richer environment. Dasgupta and Maskin [22] focused on single-valued rules and showed that majority rule uniquely satisfies anonymity, neutrality, Pareto efficiency and an independence property known as the Chernoff condition over the largest possible class of problem instances. Working on restricted domains with single-valued choice rules, Alemante et al. [1] showed that Condorcet, plurality, approval voting, and maximin rules satisfy anonymity, neutrality, and a certain monotonicity property. Conversely, Horan et al. [29] allowed for multi-valued rules and focused on the domain restriction of problems admitting strict Condorcet winners. In this setting, they characterized the rule selecting the Condorcet winner(s) with anonymity, neutrality, positive responsiveness, and an independence property they refer to as Nash independence. A further recent contribution in this line is Barberà et al. [11] (where the domain is not restricted to problems admitting strict Condorcet winners).

Regarding the literature on strategy-proof preference aggregation, Bossert and Storcken [19] were the first to study incentive-compatibility in the Arrovian setting. Working within the Kemeny framework, they established an impossibility result involving the much stronger property of group K-SP, ontoneess and a relatively esoteric invariance property of extrema independence. More recently, Bossert and Sprumont [17] proposed the notion of *betweenness* strategy-proofness (Btw-SP), according to which misreporting cannot lead to an outcome that is between the one obtained under truthful reporting and the agent’s own preferences. This property amounts to requiring that the truthful social ordering not be unambiguously dominated by the one produced under misreporting and is thus a necessary, but weak standard of non-manipulability. Bossert and Sprumont [17] identified a number of rules that satisfy Btw-SP, and axiomatized a few of them on the basis of Btw-SP and other properties. Building on these results, Athanasoglou [6] demonstrated that all rules identified in [17] violate K-SP. Harless [27] and Athanasoglou [7] investigated the interplay of Btw-SP with various solidarity properties.

Since Arrovian aggregation with Kemeny preferences admits a graph-theoretic interpretation, our work shares some parallels with the literature on single-peaked preferences and

strategyproof facility location (Moulin [39], Barberà et al. [13], Schummer and Vohra [44], Aziz et al. [8]). In these settings, so-called generalized median (or phantom) voter mechanisms are often characterized with strategyproofness and efficiency criteria. While related to voting-by-committees, these sorts of median voter mechanisms are not well-defined in the Arrovian context and thus not directly relevant.

Paper outline. The paper is organized as follows. Section 2 introduces the model. Section 3 explores the incompatibility of anonymity and neutrality, states May’s Theorem for two alternatives and defines our notions of efficiency and strategy-proofness. Section 4 drops anonymity and explores the possibilities of neutrality and our basic properties of strong unanimity and K-SP. It characterizes semi-dictator rules for two agents, and generalizes this class to arbitrary numbers of agents and alternatives while maintaining its properties. Section 5 drops neutrality and explores the possibilities of anonymity and our basic properties. For three alternatives, we define two families of rules which satisfy our basic properties and anonymity. For four alternatives and two agents, we provide the computer-aided solution for the existence of a rule satisfying our basic properties and anonymity. Section 6 provides for three alternatives and three agents a characterization of median rules with tie-breaking by strengthening strong unanimity to preference selection. Section 7 concludes. The Appendix contains all proofs omitted from the main text.

2 Model

Let $A = \{a_1, \dots, a_m\}$ denote a finite set of $m \geq 2$ alternatives and $N = \{1, \dots, n\}$ a finite set of $n \geq 2$ agents. Let \mathcal{R} denote the set of **orderings** of A (i.e. complete, reflexive, antisymmetric, and transitive binary relations). Each agent $i \in N$ has a preference relation $R_i \in \mathcal{R}$ over A . We interchangeably write aR_ib and $(a, b) \in R_i$ to denote that agent i finds alternative a at least as good as alternative b . For each $B \subset A$, let $\mathcal{R}(B)$ denote the set of orderings over B . For all $R \in \mathcal{R}$, the ordering $-R$ is defined such that for all $a, b \in A$ with $a \neq b$, $(a, b) \in R$ if and only if $(b, a) \in -R$.

A **(preference) profile** $R_N = (R_1, \dots, R_n)$ is an n -tuple of orderings, representing the preferences of all agents in N . The set of preference profiles is denoted by \mathcal{R}^N . Given $R \in \mathcal{R}$ and $B \subset A$, let $R|_B$ denote the restriction of R to B , and $R_N|_B = (R_j|_B)_{j \in N}$. If $a \in B$ and $a R_i b$ for all $b \in B$, then we say that a is the *most preferred* alternative of R_i in B .

For convenience, we often denote an ordering by listing the alternatives from left to right in increasing rank (where the first ranked alternative is the most preferred one in A). Thus, if we write $R = a_1 a_2 \dots a_m$, then alternative a_1 is ranked first, a_2 second, and so on. When

the ordering of only the first t positions of R is relevant, we write $a_1 \dots a_t \dots$ to mean that the ordering of the remaining positions can be arbitrary.

A **rule** is a function $f : \mathcal{R}^N \rightarrow \mathcal{R}$. A rule f is *dictatorial* if there exists $i \in N$ such that for all $R_N \in \mathcal{R}^N$ we have $f(R_N) = R_i$. A rule f is *constant* if $f(R_N) = f(R'_N)$ for all $R_N, R'_N \in \mathcal{R}^N$. A **choice rule** is a function $\varphi : \mathcal{R}^N \rightarrow A$.

3 Definitions and preliminary results

In this section we explore the existence of rules satisfying anonymity and neutrality. For more than two alternatives, we show for many instances we have non-existence, whereas for two alternatives with an odd number of agents we have existence by May's Theorem. Dropping either anonymity or neutrality, we would like rules to satisfy in addition efficiency and strategy-proofness notions, and define three of them for each, from weak to strong. The strongest non-manipulability notion will be Kemeny strategy-proofness, which will then be explored together with either neutrality or anonymity in two subsequent sections.

3.1 Anonymity and neutrality

How might rules incorporate a concern for fairness? Two properties, which are intuitively appealing, are anonymity and neutrality.

Anonymity requires that a rule be invariant to reshuffling agent identities. Formally, let $\sigma : N \mapsto N$ denote a permutation of N . For each profile $R_N \in \mathcal{R}^N$, let $R_{\sigma(N)} \equiv (R_{\sigma(1)}, \dots, R_{\sigma(n)})$ denote the profile where agents have been relabeled according to σ .

Anonymity. For each $R_N \in \mathcal{R}^N$ and each permutation σ of N , $f(R_N) = f(R_{\sigma(N)})$.

Thus, anonymity ensures that the identity of agents does not affect the outcome of the rule. It excludes dictatorial rules from consideration.

A second fairness property focuses on the identity of alternatives (and not agents) and requires that the rule does not systematically favor one alternative over another. Formally, let $\pi : A \mapsto A$ be a permutation of A . For each $R \in \mathcal{R}$, let $\pi R \in \mathcal{R}$ be the ordering such that for all $a, b \in A$, $\pi(a) \pi R \pi(b)$ if and only if $a R b$. For each $R_N \in \mathcal{R}^N$, let $\pi R_N \equiv (\pi R_1, \dots, \pi R_n)$.

Neutrality. For each $R_N \in \mathcal{R}^N$ and each permutation π of A ,

$$f(\pi R_N) = \pi f(R_N).$$

Neutrality ensures that all alternatives are treated equally. If it holds, the rule cannot discriminate either for or against a particular alternative. It excludes constant rules from

consideration.

While anonymity and neutrality are seemingly mild requirements, they are generally incompatible.⁵ The existence of an anonymous and neutral rule will depend on the values of n and m , the number of agents and alternatives. Theorem 1 provides a necessary and sufficient condition to this effect. The proofs of all results are relegated to the Appendix.

Theorem 1 *Let $|A| = m$ and $|N| = n$. Then there exists an anonymous and neutral rule if and only if all prime factors of n are strictly greater than m .*

Theorem 1 implies that, regardless of the number of alternatives, anonymity and neutrality are incompatible if the number of agents is even (since two is a prime factor of any even number). Thus, anonymity and neutrality are compatible for a relatively small set of problem instances. The proof of Theorem 1 is a straightforward adaptation of an earlier result by Moulin [40] that applies to choice rules (Theorem 1 of [40]). Nevertheless, it is worth noting (albeit unsurprising) that the incompatibility of neutrality and anonymity is more acute for aggregation rules than it is for choice rules. This is due to the combination of two facts (see Problem 1 of Moulin [40, p. 25]) : (1) the existence of a neutral and anonymous choice rule is equivalent to the condition that m cannot be written as the sum of the nontrivial divisors of n and (2) if m cannot be written as the sum of the nontrivial divisors of n then all prime factors of n are strictly greater than m , whereas the opposite direction does not hold.

3.2 Two alternatives: May's Theorem

Theorem 1 establishes that for the majority of problem instances, it is impossible to satisfy both anonymity and neutrality. When $m = 2$ and n is odd, however, this incompatibility becomes less stark and May's Theorem [37], a classic result in social choice, provides an important possibility result. Again, if n is even, anonymity and neutrality are incompatible regardless of m .

To state May's Theorem, we need to formally define strategy-proofness. Given $R_N \in \mathcal{R}^N$ and $R'_i \in \mathcal{R}$, the notation (R'_i, R_{-i}) denotes the profile that is identical to R_N except that the preference of agent i is equal to R'_i . Minimal strategy-proofness (Min-SP) ensures that it is impossible to obtain an outcome that is exactly identical to one's preferences, unless truthful preference revelation yields the same result.

Minimal strategy-proofness (Min-SP). There do not exist $R_N \in \mathcal{R}^N, i \in N, R'_i \in \mathcal{R}$ such that $f(R'_i, R_{-i}) = R_i \neq f(R_N)$.

⁵Note that no such incompatibility exists when rules are not *strict*, i.e. when their image is expanded from \mathcal{R} to the set of subsets of \mathcal{R} (see, e.g., Young and Levenglick [45]).

Min-SP is the absolute weakest possible standard of strategy-proofness in the preference aggregation framework. However, when $m = 2$ there is no difference between Min-SP and stronger notions of non-manipulability.⁶ The literature also studies non-manipulability by groups for two alternatives, see Barberà et al. [10] and Manjunath [36] among others. For two alternatives and an odd number of agents, the majority rule chooses for any profile the relation which is possessed by a majority of agents.

Theorem 2 (*May's Theorem [37]*). *Let $|A| = m = 2$ and $|N| = n$ be odd. Then majority rule is the only rule satisfying anonymity, neutrality and Min-SP.*

When $m = 2$ and n is even, anonymity and neutrality are incompatible by Theorem 1. The classical version of May's Theorem deals with this issue by allowing for multi-valued rules, in which case the anonymity-neutrality tradeoff vanishes.⁷ By contrast, we insist on single-valued rules and we examine how May's Theorem is modified when we drop either anonymity or neutrality. To this end, we define the following two families of rules.

Definition 1 *Let $A = \{a, b\}$. A rule f is a quota-majority rule if there exists an integer $q_a \in \{0, 1, \dots, n\}$ such that, for all $R_N \in \mathcal{R}^N$ we have $f(R_N) = ab$ if $|\{i \in N : R_i = ab\}| \geq q_a$ and $f(R_N) = ba$ otherwise.*

Definition 2 *Let $A = \{a, b\}$. A rule f is a collegial-majority rule if there exists a set $\mathcal{T} \subseteq 2^N$ satisfying*

(i) *for all $T, T' \in \mathcal{T}$ we have $T \subseteq T' \Rightarrow T = T'$ and*

(ii) *for all $T \subset N$, we have $[T_1 \subseteq T \text{ for some } T_1 \in \mathcal{T}] \Leftrightarrow [T \cap T' \neq \emptyset \text{ for all } T' \in \mathcal{T}]$,*

such that for all $R_N \in \mathcal{R}^N$ we have $f(R_N) = ab$ whenever there exists $T \in \mathcal{T}$ such that $T \subseteq \{i \in N : R_i = ab\}$ and $f(R_N) = ba$ otherwise.

Proposition 1 demonstrates how May's Theorem is modified when either neutrality or anonymity are dropped from the list of requirements a rule should satisfy.

Proposition 1 *Let $|A| = m = 2$.*

(i) (*Moulin [40]*) *Quota-majority rules are the only rules satisfying anonymity and Min-SP.*

(ii) (*Moulin [40]*) *Collegial-majority rules are the only rules satisfying neutrality and Min-SP.*

⁶For two alternatives, any choice function selecting for each profile a unique alternative corresponds to the rule choosing for this profile the strict relation where the selected alternative is preferred over the non-selected alternative.

⁷Indeed, May's Theorem is originally formulated as a characterization of majority rule on the full domain with anonymity, neutrality, and a monotonicity property referred to as *positive responsiveness*.

3.3 Efficiency and strategy-proofness

In the preference-aggregation framework, combining either anonymity or neutrality together with Min-SP imposes very weak constraints on acceptable rules (and by Theorem 1 we will not impose both anonymity and neutrality). For example, constant rules choosing always the same preference are anonymous and Min-SP, and dictatorial rules are neutral and Min-SP.

Constant rules ignore agents' preferences and violate efficiency requirements. We formulate three efficiency notions, in increasing order of strength. The first is self-explanatory.

Unanimity. For all $R_N \in \mathcal{R}^N$ and all $R \in \mathcal{R}$, if $R_i = R$ for all $i \in N$ then $f(R_N) = R$.

Our second notion of efficiency relies on the concept of betweenness.

Definition 3 Given orderings $R, R', R'' \in \mathcal{R}$, R' is **between** R and R'' , denoted by $R' \in [R, R'']$, if $R \cap R'' \subseteq R'$ (where for all $a, b \in A$, $aRb \ \& \ aR''b \Leftrightarrow (a, b) \in R \cap R''$).

It is rational to posit that if $R' \neq R''$ and $R' \in [R_i, R'']$, then agent i with preferences R_i has *unambiguous* preference for R' over R'' . For every ordering $R \in \mathcal{R}$, this binary relation on \mathcal{R} is reflexive, transitive, anti-symmetric but not complete.⁸ We refer to it as the **betweenness extension** applied to R .

Betweenness Efficiency. There do not exist $R_N \in \mathcal{R}^N$ and $R' \in \mathcal{R}$ such that $R' \in [R_i, f(R_N)]$ for all $i \in N$ and $R' \neq f(R_N)$.

Strong Unanimity. For all $R_N \in \mathcal{R}^N$, $\bigcap_{i \in N} R_i \subseteq f(R_N)$.

A rule satisfies betweenness efficiency if it selects an ordering such that there exists no other which all agents find unambiguously better. By contrast, strong unanimity applies to preference profiles in which there is unanimous agreement over individual binary comparisons. When such unanimous consensus is present, strong unanimity requires the rule to follow it. First discussed by Arrow [5], strong unanimity implies betweenness efficiency (Footnote 11 in Harless [27]) but not the other way around. A few previous papers have used the term “strong efficiency” to refer to strong unanimity [27, 7].⁹

We now address the vulnerability of a rule to strategic manipulation in a way that strengthens Min-SP. The first concept we introduce draws from the betweenness relation.

⁸It is incomplete because, given any $R \in \mathcal{R}$, there will exist multiple pairs of orderings R', R'' that satisfy $R \cap R'' \not\subseteq R'$ and $R \cap R' \not\subseteq R''$.

⁹The above efficiency requirements, together with neutrality and Min-SP, are trivially satisfied by dictatorial rules. The Online Appendix presents a non-dictatorial rule that satisfies neutrality and Min-SP but violates betweenness-efficiency and thus also strong unanimity. Hence, the efficiency requirement of strong unanimity provides a meaningful check against neutral and Min-SP rules that are very inefficient.

Betweenness strategy-proofness (Btw-SP). There do not exist $R_N \in \mathcal{R}^N, i \in N, R'_i \in \mathcal{R}$ such that $f(R'_i, R_{-i}) \in [R_i, f(R_N)]$ and $f(R'_i, R_{-i}) \neq f(R_N)$.

A rule is Btw-SP if, by misreporting one's preferences, it is not possible to obtain an outcome that is unambiguously better than the outcome under truthfulness. Evidently, Btw-SP is a nontrivial strengthening of Min-SP. As Example 1 illustrates, there do exist “unappealing” rules that satisfy neutrality, strong unanimity and Min-SP but not Btw-SP. A more complex example with anonymity instead of neutrality can be found in the Online Appendix B.

Example 1 Let $A = \{a, b, c\}$ and $|N| \geq 3$. Define the rule f as

$$f(R_N) = \begin{cases} R_1, & \text{if } R_1 = R_2 \\ R_3 & \text{otherwise.} \end{cases} \quad (1)$$

This rule is easily seen to be neutral and strongly unanimous. We now show that it is Min-SP but not Btw-SP. If $i \notin \{1, 2, 3\}$, agent i has no effect on f . If $i \in \{1, 2\}$, there exists no $R'_i \neq R_i$ such that $f(R_N) \neq R_i$ and $f(R'_i, R_{-i}) = R_i$. If $i = 3$, then $f(R_N) \neq R_i$ implies that $R_1 = R_2$. Thus $f(R'_i, R_{-i}) = R_1 = R_2$ for any deviation $R'_i \neq R_i$. So, f is Min-SP. To show that it fails Btw-SP consider the profile $(R_1, R_2, R_3) = (abc, bac, cba)$. We have $f(R_N) = cba$. Let $R'_1 = bac$. Then $f(R'_1, R_2, R_3) = bac \in [abc, cba]$, which results in a violation of Btw-SP.

Beginning with the work of Bossert and Sprumont [17], various rules have been found to satisfy Btw-SP, and it has formed the basis of various characterizations [17, 7, 27]. Though stronger than Min-SP, this property still provides a relatively weak notion of non-manipulability.

While Btw-SP is a useful benchmark for strategy-proofness, the incompleteness of the betweenness relation diminishes its impact. Indeed, if two orderings are incomparable, then we cannot say whether a preference misreport is profitable or not. Therefore, we search for a way to capture preferences over orderings that is consistent with betweenness when the latter produces clear results, but that also yields a complete relation. To this end, we follow a two-stage approach. First, we determine a way to measure the distance between two orderings. Second, we employ this concept of distance to propose a way of ranking orderings in \mathcal{R} .

To guide the first part of our exercise, we require three basic requirements that a distance function on \mathcal{R} should satisfy: (i) metric conditions, (ii) consistency with betweenness¹⁰;

¹⁰In the sense that if $R' \in [R, R'']$, then the distance from R to R'' is equal to the sum of the distance or R to R' and the distance of R' to R'' .

and (iii) invariance to relabeling of the alternatives. Improving on the classic result of Kemeny and Snell [31], Can and Storcken [21] showed that properties (i)-(ii)-(iii) together with a normalization requirement that the minimal non-zero distance equals one uniquely characterize the Kemeny distance [30], a well-known metric in the space of orderings. For completeness, we provide their characterization below.¹¹ The formal definition follows.

Definition 4 *Given two orderings $R, R' \in \mathcal{R}$, let $D(R, R') = (R \setminus R') \cup (R' \setminus R)$. The **Kemeny distance** between R and R' , denoted by $\delta(R, R')$, is defined as $\delta(R, R') = \frac{|D(R, R')|}{2}$.*

In words, $\delta(R, R')$ is the number of alternative pairs on whose relative ranking the two orderings disagree. For example, if $R = abc$ and $R' = cab$, then $R \setminus R' = \{(a, c), (b, c)\}$, $R' \setminus R = \{(c, a), (c, b)\}$ and $\delta(R, R') = 2$.

Theorem 3 [Can and Storcken (2018)] *Let function $\alpha : \mathcal{R} \times \mathcal{R} \mapsto \mathbb{R}$ denote the distance between two orderings. Then α is the Kemeny distance if and only if it satisfies the following four conditions:*

Condition 1 (Metric conditions). *For all $R, R', R'' \in \mathcal{R}$, we have:*

- (i) *Non-negativity: $\alpha(R, R') \geq 0$.*
- (ii) *Identity of indiscernibles: $\alpha(R, R') = 0$ if and only if $R = R'$.*
- (iii) *Symmetry: $\alpha(R, R') = \alpha(R', R)$.*
- (iv) *Triangle inequality: $\alpha(R, R'') \leq \alpha(R, R') + \alpha(R', R'')$.*

Condition 2 (Betweenness). *For all $R, R', R'' \in \mathcal{R}$ such that $R' \in [R, R'']$ we have $\alpha(R, R'') = \alpha(R, R') + \alpha(R', R'')$.*

Condition 3 (Neutrality). *For all $R, R' \in \mathcal{R}$ and all permutations π on A , we have $\alpha(R, R') = \alpha(\pi R, \pi R')$.*

Condition 4 (Normalization). $\min\{\alpha(R, R') : R \neq R'\} = 1$.

¹¹In a recent contribution, Nishimura and Ok [41] proposed and axiomatized an alternative class of semimetrics for acyclic preference relations. In contrast to the Kemeny metric, these semimetrics are directly motivated by their implications for choice behavior. The most prominent member of this class is the “top difference semimetric,” which uniquely has the added feature of being interpretable as a weighted version of the Kemeny metric (instances of which were studied in Can [20] and Hassanzadeh and Milenkovic [28]).

With Definition 4 in mind, every ordering $R \in \mathcal{R}$ induces a complete, reflexive and transitive binary relation on \mathcal{R} , the **Kemeny extension** applied to R , whereby orderings are ranked on the basis of their Kemeny distance to R . The smaller this distance, the more preferred is the ordering. This binary relation stipulates that two Kemeny-equidistant orderings from R are indifferent for an agent with preference R .¹²

Given any $R \in \mathcal{R}$, it is easy to verify that for all $R', R'' \in \mathcal{R}$ such that $R' \neq R''$, if $R' \in [R, R'']$, then $\delta(R, R') < \delta(R, R'')$. Thus, the Kemeny extension of a preference preserves the betweenness relation.¹³

We now introduce the incentive-compatibility property based on the Kemeny extension.

Kemeny strategy-proofness (K-SP). There do not exist $R_N \in \mathcal{R}^N, i \in N, R'_i \in \mathcal{R}$ such that $\delta(R_i, f(R'_i, R_{-i})) < \delta(R_i, f(R_N))$.

A rule is K-SP if no misrepresentation yields an outcome which is more preferred according to the Kemeny extension (applied to the deviating agent’s preference) than the one obtained under truthfulness.¹⁴ In other words, K-SP ensures that by misreporting, no agent can obtain an outcome that is closer to his true preference according to the Kemeny distance.

Since the Kemeny extension preserves the betweenness extension, K-SP implies Btw-SP. In fact, it strengthens the latter property significantly: none of the known nontrivial Btw-SP rules satisfy it, unless the number of alternatives is restricted to three (see Athanassoglou [6] and Section 4). Along related lines, Bossert and Storcken [19] established an impossibility result involving the much stronger coalitional version of K-SP, ontoneess, and a relatively esoteric invariance property to which they refer as extrema independence.

Verifying whether K-SP is satisfied by a given rule can be difficult, as comparing orderings on the basis of their Kemeny distance from a certain benchmark is not easy. Computer simulations are often needed to generate counterexamples, even for problem instances of small size [6]. For this reason, when investigating a rule’s K-SP, it would be helpful to restrict the set of preference misrepresentations that need to be compared to truthful reporting.

One way of achieving this goal is by showing that small deviations from truthfulness are sufficient to adjudicate the rule’s K-SP. Along these lines, a number of recent papers have focused on identifying necessary and sufficient conditions for the equivalence between global

¹²Our restriction of the set of preferences over orders, although natural, may not be the only possible one. Other ways to obtain preferences over orders from a “true” order appear in Amorós, Corchón, and Moreno [3] and Amorós [2] among others.

¹³Despite its simplicity and desirable properties, the Kemeny extension can sometimes yield implications that are at odds with menu-based choice behavior (see Ex. 1.1 in Nishimura and Ok [41] for a simple illustration). We should thus be aware of its limitations as we use it to define a stronger standard of strategy-proofness.

¹⁴In what follows, and in a slight abuse of grammar, we use the acronym K-SP to denote both “Kemeny strategy-proofness” as well as “Kemeny strategy-proof”.

and local measures of strategy-proofness [42, 33, 34]. Since the results of those papers do not readily apply to the Arrovian aggregation framework with Kemeny-based preferences, we explore the local-global equivalence directly.

Before proceeding, we specify what we mean by local measures of strategy-proofness in our setting.

Local Kemeny strategyproofness (Local K-SP). There do not exist $R_N \in \mathcal{R}^N, i \in N, R'_i \in \mathcal{R}$ such that $\delta(R_i, R'_i) = 1$ and $\delta(R_i, f(R'_i, R_{-i})) < \delta(R_i, f(R_N))$.

Thus, a rule is Local K-SP if by misreporting the order of a unique adjacent alternative pair, it is not possible to obtain an outcome that is closer in Kemeny distance to one's true preferences. Clearly, K-SP implies Local K-SP. The following result establishes that the opposite holds as well, provided the rule also satisfies Min-SP, the weakest possible measure of global non-manipulability. Note that Local K-SP does not imply Min-SP.¹⁵

Proposition 2 *If a rule satisfies Local K-SP and Min-SP, then it satisfies K-SP.*

Proposition 2 will be a useful tool in establishing the K-SP of the rules we introduce in the next section as we only have to show Min-SP and K-SP for deviations where the ordering of a unique adjacent alternative pair is reversed (instead of any deviation).

4 Keep neutrality - Drop anonymity

In this section we explore strongly unanimous and K-SP rules that satisfy neutrality but fail anonymity. We are able to establish a full characterization when the number of agents is two.

4.1 Two agents

We focus on the two-agent case and define a family of rules that forms the cornerstone of this section.

Definition 5 *A two-agent semi-dictator rule is parameterized by the following two inputs:*

- (i) A **semi-dictator** $i \in N = \{1, 2\}$.

¹⁵We provide an example in the Online Appendix. We conjecture, but so far have been unable to prove, that local K-SP combined with some mild efficiency property implies Min-SP. This would mean that, given mild efficiency requirements, local K-SP is equivalent to K-SP. Such a result would more closely align with the contribution of Kumar et al. [34].

(ii) A **position set** $P \subset \{1, 2, \dots, m-1\}$ satisfying for all distinct $p, p' \in P$, $|p - p'| > 2$.

Let $R_N \in \mathcal{R}^N$. Without loss of generality, suppose that the semi-dictator i has preference $R_i = a_1 a_2 \dots a_m$. The semi-dictator rule chooses for R_N the ordering $f^{(i,P)}(R_N)$ whose k -th alternative is defined as follows: for all $k \in \{1, 2, \dots, m\}$,

$$f_k^{(i,P)}(R_N) = \begin{cases} a_{k+1}, & \text{if } k \in P \text{ and } a_{k+1} R_j a_k \\ a_{k-1}, & \text{if } k-1 \in P \text{ and } a_k R_j a_{k-1} \\ a_k, & \text{otherwise.} \end{cases} \quad (2)$$

where $N = \{i, j\}$.

A two-agent semi-dictator rule $f^{(i,P)}$ produces an ordering that is identical to the preferences of the semi-dictator i except possibly at the alternatives occupying ranks $\{p, p+1\}$ where $p \in P$. In particular, given the semi-dictator's preferences $R_i = a_1 a_2 \dots a_m$, for every position $p \in P$, alternatives a_p and a_{p+1} will be assigned rank either p or $p+1$, in accordance with agent j 's preferences.

Figure 1 illustrates a two-agent semi-dictator rule when $m = 14$, semi-dictator 1, $P = \{4, 7, 13\}$ and $R_1 = a_1 a_2 \dots a_{14}$.




Figure 1: An illustration of a two-agent semi-dictator rule $f = f^{(i,P)}$ with $i = 1$, $P = \{4, 7, 13\}$ and $R_1 = a_1 a_2 \dots a_{14}$. The rule applied to profile R_N produces a social ordering that is identical to R_i , except possibly at ranks (4,5), (7,8) and (13,14) where the relative order of adjacent alternatives is determined by agent 2's preferences R_2 . For example, if agent 2 prefers a_5 to a_4 , a_7 to a_8 and a_{14} to a_{13} , then $f(R_N) = a_1 a_2 a_3 [a_5 a_4] a_6 [a_7 a_8] a_9 a_{10} a_{11} a_{12} [a_{14} a_{13}]$, where the brackets are added for emphasis.

It is important that any two distinct positions in the set P have distance greater than two as otherwise the two-agent semi-dictator rule might be manipulated by the semi-dictator. We illustrate this below for five alternatives where either $P = \{1, 3\}$ or $P = \{1, 4\}$.

Example 2 Let $A = \{a_1, a_2, a_3, a_4, a_5\}$ and $N = \{1, 2\}$.

On the one hand, if $i = 1$ and $P = \{1, 3\}$, then the semi-dictator rule is not K-SP: let $R_1 = a_1a_2a_3a_4a_5$ and $R_2 = a_2a_1a_4a_3a_5$; then $f^{(1,P)}(R_1, R_2) = a_2a_1a_4a_3a_5$ and $\delta(R_1, f^{(1,P)}(R_1, R_2)) = 2$; for $R'_1 = a_1a_3a_2a_4a_5$ we have $f^{(1,P)}(R'_1, R_2) = R'_1$ and $\delta(R_1, R'_1) = 1$, and agent 1 profitably manipulates from (R_1, R_2) via R'_1 .

On the other hand, if $i = 1$ and $P = \{1, 4\}$, then the semi-dictator rule is K-SP. Consider first agent 1. Let $R_1 = a_1a_2a_3a_4a_5$ and R_2 be arbitrary. Then by definition, $\delta(R_1, f^{(1,P)}(R_1, R_2)) \leq 2$. If $\delta(R_1, f^{(1,P)}(R_1, R_2)) \leq 1$, then it is easy to see that agent 1 cannot manipulate. Thus, let $\delta(R_1, f^{(1,P)}(R_1, R_2)) = 2$. Now agent 1 cannot manipulate by pushing a_1 or a_2 to fourth or fifth position in his deviation R'_1 as then $\delta(R_1, f^{(1,P)}(R'_1, R_2)) \geq 2$. Similarly agent 1 cannot manipulate by pushing a_4 or a_5 to first or second position in his deviation. Now in the deviation R'_1 either the first two ranked alternatives are $\{a_1, a_2\}$ or the last two ranked alternatives are $\{a_4, a_5\}$. Then when a_3 is ranked third we have $f^{(1,P)}(R'_1, R_2) = f^{(1,P)}(R_1, R_2)$, when a_3 is ranked first or second we have $f^{(1,P)}(R'_1, R_2)|_{\{a_4, a_5\}} = f^{(1,P)}(R_1, R_2)|_{\{a_4, a_5\}}$ and $\delta(R_1, f^{(1,P)}(R'_1, R_2)) \geq 2$, and when a_3 is ranked fourth or fifth we have $f^{(1,P)}(R'_1, R_2)|_{\{a_1, a_2\}} = f^{(1,P)}(R_1, R_2)|_{\{a_1, a_2\}}$ and $\delta(R_1, f^{(1,P)}(R'_1, R_2)) \geq 2$. Now consider agent 2. By misreporting his preferences, this agent can at most change the relative order of a_1 and a_2 at the top of the ordering or the relative order of a_4 and a_5 at the bottom of the ordering, or both. Since these relative orders are, by definition, consistent with agent 2's preferences, we may immediately deduce that $\delta(R_2, f(R_1, R_2)) < \delta(R_2, f(R_1, R'_2))$ for any R'_2 such that $f(R_1, R_2) \neq f(R_1, R'_2)$. The detailed argument can be found in the proof of Theorem 5.

Our first main contribution is the characterization of two-agent semi-dictator rules with strong unanimity, neutrality and K-SP.

Theorem 4 Let $N = \{1, 2\}$ and $|A| = m \geq 2$. A rule f satisfies strong unanimity, neutrality and K-SP if and only if f is a two-agent semi-dictator rule.

The proof of Theorem 4 proceeds by establishing the characterization for $m = 3$ and $m = 4$, and then tackles the case $m \geq 4$ by induction. The argument for $m = 4$ is particularly involved as it requires the careful examination of many different sub-cases.

For two alternatives, if both agents report the same preference, then by strong unanimity this preference is chosen by the rule; and if the two agents disagree on the ranking of the two

alternatives, then by neutrality always the same agent's ranking is chosen for all profiles, i.e. the rule is semi-dictatorial with empty position set (and K-SP is redundant).¹⁶

4.2 More Than Two Agents

Our second main contribution is to define the general class of semi-dictator rules and to show that they satisfy the properties of Theorem 4. In doing so, we also identify the first nontrivial K-SP rule for an arbitrary number of agents and alternatives. Unlike the two-agent case, a characterization of semi-dictator rules for general n and m remains elusive.

We begin by defining the concept of a committee, which plays a central role in the analysis. Committees and extensions thereof (e.g., left-right coalition systems) have been studied extensively in a variety of models of social choice (Barberà [9]).

Definition 6 *A **committee** is a non-empty collection \mathcal{C} of subsets of N satisfying the following two conditions:*

- (1) *for all $C, C' \subset N$, we have $C \in \mathcal{C} \& C \subset C' \Rightarrow C' \in \mathcal{C}$ and*
- (2) *for all $C \subset N$, we have $C \in \mathcal{C} \Leftrightarrow N \setminus C \notin \mathcal{C}$.*

Committees serve the following function in semi-dictator rules, which is reminiscent to the “voting by committee” procedure of Barberà et al. [14]. Given any pair $\{a, b\} \subset A$, a committee \mathcal{C} indicates the set of the so-called “winning coalitions” when deciding the order of alternatives a and b . In particular, if the preference profile in question is such that the set of agents preferring a to b is contained in \mathcal{C} , then the semi-dictator rule will rank a before b . This procedure is well-defined since agents have complete preferences over alternatives and the definition of committees ensures that $C \in \mathcal{C}$ iff $N \setminus C \notin \mathcal{C}$.

Clearly, when $m = 2$, the voting by committees procedure is K-SP. The challenge is to design an aggregation rule that integrates voting by committee when $m \geq 3$ in a way that does not lead to violations of transitivity and K-SP. This is exactly what semi-dictator rules accomplish.

Definition 7 *A **semi-dictator rule** is parameterized by the following three inputs:*

- (i) *A **semi-dictator** $i \in N$.*
- (ii) *A **position set** $P \subset \{1, 2, \dots, m - 1\}$ satisfying for all distinct $p, p' \in P$, $|p - p'| > 2$.*

¹⁶We are grateful to Reviewer 2 for pointing this out to us for two alternatives.

(iii) For each position $p \in P$ a **committee** \mathcal{C}_p on $N \setminus \{i\}$. Let $\mathcal{C} \equiv \mathcal{C}_p|_{p \in P}$ denote the corresponding family of committees.

Let $R_N \in \mathcal{R}^N$. Without loss of generality, suppose that the semi-dictator i has preferences $R_i = a_1 a_2 \dots a_m$. The semi-dictator rule chooses for R_N the ordering $f^{(i,P,\mathcal{C})}(R_N)$ whose k -th alternative is defined as follows: for all $k \in \{1, \dots, m\}$,

$$f_k^{(i,P,\mathcal{C})}(R_N) = \begin{cases} a_{k+1}, & \text{if } k \in P \text{ and } \{j \in N \setminus \{i\} : a_{k+1} R_j a_k\} \in \mathcal{C}_k \\ a_{k-1}, & \text{if } k-1 \in P \text{ and } \{j \in N \setminus \{i\} : a_k R_j a_{k-1}\} \in \mathcal{C}_{k-1} \\ a_k, & \text{otherwise.} \end{cases} \quad (3)$$

A semi-dictator rule $f^{(i,P,\mathcal{C})}$ when applied to a profile R_N produces an ordering that is identical to the preferences of the semi-dictator i except possibly at the alternatives occupying ranks $\{p, p+1\}$ where $p \in P$. Given the semi-dictator's preferences $R_i = a_1 a_2 \dots a_m$, for every position $p \in P$, alternatives a_p and a_{p+1} will be assigned rank either p or $p+1$. If the set of agents preferring a_p to a_{p+1} in profile R_N (i.e., the set $\{j \in N \setminus \{i\} : a_p R_j a_{p+1}\}$) belongs to the committee \mathcal{C}_p , then alternative a_p is assigned rank p and a_{p+1} rank $p+1$, consistent with the semi-dictator's preferences; if not, a_{p+1} is assigned rank p and a_p rank $p+1$, in contrast to the semi-dictator's preference. This procedure is well-defined because there is always a gap between pairs of adjacent alternatives whose order is decided by committee – this is guaranteed by the requirement that if $p, p' \in P$ such that $p \neq p'$, then $|p - p'| > 2$. Furthermore, analogous to Example 2 it can be seen that the gap has to be greater than two as otherwise the semi-dictator might be able to manipulate the rule.

Figure 2 illustrates a semi-dictator rule when $m = 14$, semi-dictator 1 with $R_1 = a_1 a_2 \dots a_{14}$ and $P = \{4, 7, 13\}$. The structure of the committees $\mathcal{C}_4, \mathcal{C}_7, \mathcal{C}_{13}$ is specified in the figure and its caption.

While semi-dictator rules are neutral, they are obviously not anonymous. A way of improving their fairness from the point of view of the agents is by maximizing the number of alternative pairs to be decided by committees. Along these lines, the number of alternatives pairs whose relative order is decided by committee can range from 0 (when the semi-dictator is in fact a dictator) to $\lfloor \frac{m+1}{3} \rfloor$. If we wanted to constrain the semi-dictator's power as much as possible ex-ante, we would choose a semi-dictator rule with $|P| = \lfloor \frac{m+1}{3} \rfloor$.

We now turn to the efficiency and incentive properties of semi-dictator rules. The main result we establish is the K-SP of all semi-dictator rules. As mentioned earlier, Proposition 2 allows us to simplify the proof by focusing only on adjacent deviations from truthful reporting.

Theorem 5 *Semi-dictator rules satisfy strong unanimity, neutrality and K-SP.*




Figure 2: An illustration of a semi-dictator rule $f = f^{(i,P,C)}$ with $i = 1$, $P = \{4, 7, 13\}$ and $R_1 = a_1 a_2 \dots a_{14}$. The rule applied to profile R_N produces a social ordering that is identical to R_1 , except possibly at ranks (4,5), (7,8) and (13,14) where the relative order of adjacent alternatives: (i) $\{a_4, a_5\}$ is determined by majority rule with ties broken against the semi-dictator; (ii) $\{a_7, a_8\}$ follows the semi-dictator's wishes unless all other agents rank a_8 before a_7 ; and (iii) $\{a_{13}, a_{14}\}$ goes against the semi-dictator's wishes as long as both agents 2 and 4 prefer a_{14} to a_{13} .

Remark 1 *It is worth noting that semi-dictator rules can be generalized to allow for committees that depend not only on the position set P , but also on the alternative pairs whose order the committee determines. In other words, we could define semi-dictator rules where we introduce for each position $p \in P$ and unordered pair of alternatives $\{a, b\} \subset A$ a **committee** $C_p(\{a, b\})$. The proof of K-SP, detailed in Theorem 5, carries over to this more general setting. Of course, if we extend semi-dictator rules in this manner they will fail to be neutral. Furthermore, for three agents and three alternatives we later establish in Proposition 4 that rules other than semi-dictator rules satisfy strong unanimity, neutrality and K-SP.*

5 Keep anonymity - Drop neutrality

In this section we explore strongly unanimous and K-SP rules that satisfy anonymity but fail neutrality. In contrast to the previous sections, we are not able to find a family of rules that satisfies these properties on the full domain. Instead, we establish possibility results for the two special cases where there are three alternatives or there are four alternatives and two agents. We suspect, but have been unable to prove, that there exists no strongly unanimous,

anonymous and K-SP rule for general n and m .

5.1 Three alternatives

We begin by defining a set of orderings of the elements of \mathcal{R} that will prove useful later on. Note that for all $R \in \mathcal{R}$, the ordering $-R$ is defined so that for all $a, b \in A$ such that $a \neq b$, $(a, b) \in R$ if and only if $(b, a) \in -R$. Since we drop neutrality, the rules that will be introduced in this section treat some alternatives (i.e. orderings) more favorably than others. Specifically, the rules are based on an exogenous ordering of the alternatives, which we denote by \succeq .

Definition 8 *An ordering \succeq of \mathcal{R} is **regular** if, for all $R \in \mathcal{R}$, whenever R is ranked first by \succeq , then for all R', R'' different than R and $-R$ and such that $R'' \in [R', R]$, we cannot have both $-R \succeq R'$ and $-R \succeq R''$.*

For example, if \succeq is regular and ranks abc first, then we cannot have both $cba \succeq bac$ and $cba \succeq bca$, and we also cannot have both $cba \succeq acb$ and $cba \succeq cab$. An example of an ordering \succeq that is regular is one that ranks orderings on the basis of their Kemeny distance from a benchmark R (the smaller the distance, the higher the rank), with ties broken arbitrarily. We may call such an ordering *Kemeny-consistent*.

We proceed by defining two families of rules that are known in the literature and play an important role in this section.

Definition 9 *Let \succeq be an ordering on \mathcal{R} . For all $R_N \in \mathcal{R}^N$, let*

$$K(R_N) = \arg \min_{R \in \mathcal{R}} \sum_{i \in N} \delta(R, R_i). \quad (4)$$

The \succeq -Condorcet-Kemeny rule is defined as the aggregation rule which assigns to each $R_N \in \mathcal{R}^N$ the ordering belonging to $K(R_N)$ ranked highest according to \succeq .

Definition 10 *Let \succeq be an ordering on \mathcal{R} . Rule f is the **fixed-benchmark rule**¹⁷ associated with \succeq if, for all $R_N \in \mathcal{R}^N$,*

$$f(R_N) = R \text{ where } R \supseteq \bigcap_{i \in N} R_i \text{ and } R \succeq R' \text{ for all } R' \in \mathcal{R} \text{ such that } R' \supseteq \bigcap_{i \in N} R_i. \quad (5)$$

¹⁷The literature refers to those rules as fixed-order status-quo rules. As we will use the term of status-quo later, we will refer to them here as “fixed-benchmark rules”.

Condorcet-Kemeny and fixed-benchmark rules are strongly unanimous, anonymous and Btw-SP [17, 7]. Since they use an exogenous ordering \succeq on \mathcal{R} to break ties, they violate neutrality.¹⁸

Proposition 3 demonstrates that, when $m = 3$, any \succeq -Condorcet-Kemeny and any \succeq -fixed-benchmark rule will satisfy K-SP if and only if the ordering \succeq is regular.

Proposition 3 *Let $|A| = m = 3$.*

- (1) *The \succeq -Condorcet-Kemeny rule satisfies K-SP if and only if \succeq is regular.*
- (2) *The \succeq -fixed-benchmark rule satisfies K-SP if and only if \succeq is regular.*

Unfortunately, Proposition 3 does not extend to four or more alternatives. This was already known for Condorcet-Kemeny rules, as Athanasoglou [6] showed that all such rules will fail K-SP for $m \geq 4$ and $n \geq 5$. As for fixed-benchmark rules, we show why all of them will fail K-SP for $m \geq 4$ and $n = 12$. Without loss of generality, suppose $m = 4$ (as for $m > 4$ we let all agents rank $m - 4$ alternatives at the bottom identically) and suppose f is a \succeq -fixed-benchmark rule such that \succeq ranks $abcd$ first. Consider the profile R_N with 12 agents where each agent has a different ordering and $\bigcap_{i \in N} R_i = (d, a)$.¹⁹ Then there exists exactly one agent j such that $\delta(R_j, f(R_N)) = 5$, e.g., if $f(R_N) = dabc$, then this agent j has preference $R_j = cbda$. Now, if agent j deviates to $R'_j = abcd$, then $f(R'_j, R_{-j}) = abcd$ and $\delta(R_j, abcd) = 4$, a violation of K-SP. A similar argument works for any other \succeq -fixed-benchmark rule.

5.2 Four alternatives and two agents

Below we focus on the case of four alternatives and two agents.

We begin by showing that even in such environments, both families of rules considered in Proposition 3 fail K-SP. We restrict attention to Kemeny-consistent orderings \succeq but suspect a similar reasoning to hold for any other \succeq that are regular without being Kemeny-consistent. Suppose f is a \succeq -Condorcet-Kemeny rule or a fixed-benchmark rule where \succeq is Kemeny-consistent with $abcd$ as the highest-ranked ordering (as we will see both rules yield identical outcomes in the following example). Consider the profile $(R_1, R_2) = (cbda, dabc)$,

¹⁸To be precise, fixed-benchmark rules are parameterized with a special kind of *partial order* on \mathcal{R} that is referred to as *conclusive* (Athanasoglou [7]). To avoid uninteresting complications, we focus on fixed-benchmark rules which employ a full linear ordering \succeq .

¹⁹Note that there are 6 orderings ranking d first, there are 4 orderings ranking d second and above a , and there are 2 orderings ranking d third and above a .

and suppose $dabc \succeq bcda$ so that $f(R_1, R_2) = dabc$ ²⁰ and $\delta(R_1, f(R_1, R_2)) = 5$. Then the deviation $R'_1 = cbad$ yields $f(R'_1, R_2) = abcd$, leading to $\delta(R_1, f(R'_1, R_2)) = 4$ and a violation of K-SP. Relabeling alternatives, we conclude that for all Kemeny-consistent orderings \succeq we can construct a two-agent problem where K-SP is violated for both types of rules.

The failure of the rules of Proposition 3 means that we have to search elsewhere for possible strongly unanimous, anonymous and K-SP rules.

Theorem 6 *Let $N = \{1, 2\}$ and $|A| = m = 4$. There exists a rule satisfying strong unanimity, anonymity and K-SP.*

We established Theorem 6 by framing the existence of an anonymous, strongly unanimous and K-SP rule as an *integer program* and obtained a computational solution in Matlab. All details are available in the Online Appendix in which the integer program and its implementation in Matlab are described. An Excel file containing the output of all 576 profiles is included in the online Supplementary Material.

The calculated family of rules satisfying the desired properties have the following characteristics:

1. A *losing alternative* (say a) is identified and placed as low as possible in the society's ranking subject to respecting strong unanimity. For example, for any $R \in \mathcal{R}$, $f(R, -R)$ always places the losing alternative a at the bottom.
2. All other alternatives are treated symmetrically in the sense that they have identical rank-frequency vectors as detailed below.

If we evaluate the rule at all possible $((4!)^2 = 576)$ profiles, we obtain the following rank-frequency matrix:

	1	2	3	4
a	36	84	156	300
b	180	164	140	92
c	180	164	140	92
d	180	164	140	92

Table 1: Cell $[x, k]$ indicates the number of profiles in which alternative x is ranked k th by the rule. Here alternative a is the losing alternative.

The calculated rule has the following features. When both agents rank the same alternative at the bottom, then by strong unanimity this alternative is ranked at the bottom by the social

²⁰If $bcda \succeq dabc$, then we repeat a similar reasoning for the profile $(dach, bcda)$.

ordering. Now when considering the subdomain where both agents rank the same non-losing alternative at the bottom, say d , then the rule restricted to the other three alternatives is a rule satisfying strong unanimity, anonymity and K-SP. It turns out that those rules are “fixed-status-quo rules with tie-breaking”. This also applies to the subdomains where both agents rank the same non-losing alternative at the top (and by strong unanimity this alternative is at the top of the social ordering). We define below such rules.

Example 3 Let $N = \{1, 2\}$ and $A = \{a, b, c\}$. Fix a status-quo ordering R_0 , say $R_0 = cba$. The R_0 -fixed-status-quo rule with tie-breaking f makes the following choice for any profile $R_N = (R_1, R_2) \in \mathcal{R}^N$:

- (i) if $\bigcap_{i \in N} R_i \subseteq R_0$, then $f(R_N) = R_0$;
- (ii) if $\bigcap_{i \in N} R_i \not\subseteq R_0$ and $R_1 \in [R_0, R_2]$, then $f(R_N) = R_1$;
- (iii) if $\bigcap_{i \in N} R_i \not\subseteq R_0$ and $R_2 \in [R_0, R_1]$, then $f(R_N) = R_2$; and
- (iv) otherwise, we have either $[R_1 = acb \& R_2 = bac]$ or $[R_1 = bac \& R_2 = acb]$ and set $f(R_N) = bac$.²¹

It is obvious that f satisfies strong unanimity and anonymity, and one can also check K-SP.

Indeed, the rule described in Example 3 is the fixed-status-quo rule with tie-breaking when both agents rank the non-losing alternative d at the bottom. Let \underline{R}_0^d denote the fixed-status-quo ordering when both agents rank d at the bottom, and \overline{R}_0^d denote the fixed-status-quo ordering when both agents rank d at the top. Then, for the rule we found, it holds that $\underline{R}_0^d = cba$ and $\overline{R}_0^d = bca$, i.e. a is ranked as low as possible by \underline{R}_0^d and \overline{R}_0^d and the order of b and c is reversed for those two fixed-status-quo orderings. This pattern is confirmed as it holds for any non-losing alternative and we detail all the fixed-status-quo orderings when both agents rank a non-losing alternative at the bottom or at the top (see also Figure 3).

$$\begin{aligned} \underline{R}_0^b &= dca & \overline{R}_0^b &= cda \\ \underline{R}_0^c &= bda & \overline{R}_0^c &= dba \\ \underline{R}_0^d &= cba & \overline{R}_0^d &= bca \end{aligned} \tag{6}$$

Note also the following: the non-losing alternatives are ranked in a cycle by the fixed-status-quo orderings where a non-losing alternative is ranked at the bottom as \underline{R}_0^b ranks d before c ,

²¹More precisely, R_i is chosen if the third ranked alternative of R_i and the first ranked alternative of R_0 coincide.




Figure 3: Fixed-status-quo orderings (indicated in capital letters) when both agents rank a non-losing alternative at the bottom or at the top. Any two adjacent orderings in the hexagon differ by switching the ordering of two adjacent alternatives. For the first hexagon, both agents rank alternative b at the bottom. The numbers refer to the Excel file and the column FixBottom/FixTopProfiles.




Figure 4: Rotating-status-quo rules with tie-breaking where both agents rank a at the bottom or at the top. The outcomes are indicated in capital letters while italic letters are used for preferences, and the numbers refer to the table in the online Excel file.

\underline{R}_0^c ranks b before d , and \underline{R}_0^d ranks c before b . Now the reverse holds for the fixed-status-quo orderings where a non-losing alternative is ranked at the top.

Conversely, a “rotating-status-quo rule with tie-breaking” appears when both agents rank the losing alternative a at the bottom. Then the rule makes the following choices:

$$f(bcda, dcba) = cdba, \quad f(bdca, cdba) = dbca \quad \text{and} \quad f(cbda, dbca) = bcda. \quad (7)$$

Now the same applies to the subdomain where both agents rank the losing alternative at the top (and by strong unanimity this alternative is at the top of the social ordering). That is, we again have a “rotating-status-quo rule with tie-breaking”, with the difference that the rotating status quo is the opposite of the one where the losing alternative is ranked at the bottom, i.e.

$$f(abcd, adcb) = abdc, \quad f(abdc, acdb) = acbd \quad \text{and} \quad f(acbd, adbc) = adcb. \quad (8)$$

We indicate the outputs of (7) and (8) in Figure 4. The output of the rule and its complete formal definition are given in an Excel file contained in the Online Appendix (but can also

be derived from the previous results using strong unanimity and K-SP).²² We also give the formal definition of the rule below. For any profile $R_N \in \mathcal{R}^N$,

- if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \emptyset$, then $f(R) = f(R_1|_{\{b, c, d\}}a, R_2|_{\{b, c, d\}}a)$ (i.e. the outcome of the rule is the same as for the profile where a is pushed to the bottom for both agents and the rotating-status-quo rule with tie-breaking given by (7) is used);
- if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \{(a, x)\}$ (where $x \in \{b, c, d\}$), then $f(R) = f(R_1|_{A \setminus \{x\}}x, R_2|_{A \setminus \{x\}}x)$ (i.e. the outcome of the rule is the same as for the profile where x is pushed to the bottom for both agents and the \underline{R}_0^x -fixed-status-quo rule given by (6) is used);
- if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \{(a, y), (a, z)\}$ (where $\{b, c, d\} = \{x, y, z\}$), then $f(R) = f(xR_1|_{A \setminus \{x\}}, xR_2|_{A \setminus \{x\}})$ (i.e. the outcome of the rule is the same as for the profile where x is pushed to the top for both agents and the \overline{R}_0^x -fixed-status-quo rule given by (6) is used); and
- if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \{(a, b), (a, c), (a, d)\}$, then both agents rank a first and $f(R)$ is the outcome of the rotating-status-quo rule with tie-breaking given by (8) where both agents rank a first.

The rule f satisfies strong unanimity, anonymity and K-SP (for two agents and four alternatives).

6 Preference selection

In May's theorem, the rule always chooses a preference of one of the agents. The same holds for the median voter rule choosing always the peak of one of the agents (and by Black's [15] theorem in symmetric single-peaked environments the median voter's preference is the majority relation). The property below adapts the one of "peak selection", often used in problems with a single public good, to our context.

Preference selection: For all $R_N \in \mathcal{R}^N$, we have $f(R_N) \in \{R_1, \dots, R_n\}$.

²²For instance, if $R_1 = dcba$, then (i) for $R_2 = dbca$ by strong unanimity we have $f(dcba, dbca) \in \{dbca, dcba\}$, and by K-SP and $f(bdca, cdba) = dbca$ we obtain $f(dbca, dcba) = dbca$, (ii) for $R_2 = bdca$ we have $f(dcba, bdca) = dbca$ from K-SP and $f(bdca, cdba) = dbca$, and (iii) for any $R_2 \neq bdca, dbca$, $f(R_1, R_2)$ is determined by K-SP and $f(dcba, bcda) = cdba$.




Figure 5: For three alternatives, the preferences abc , cab , and bca are K -equidistant from each other as $\delta(abc, cab) = \delta(abc, bca) = \delta(cab, bca) = 2$, and similarly for acb , bac , and cba . Profiles are triangular (or belong to \triangle) if the three reported preferences are K -equidistant from each other.

Preference selection also corresponds to the fact that always a member of the society shall be chosen to represent the social preference. Again this is in the vein of macroeconomics where the representative consumer shall be a member of the society. Note that preference selection implies strong unanimity and betweenness efficiency.

On the one hand, preference selection, neutrality and K -SP are compatible as dictatorial rules satisfy all these properties.

On the other hand, we present two impossibility results for preference selection in conjunction with anonymity and K -SP. In other words, if always some agent is chosen to represent society, then either anonymity or K -SP is violated.

Theorem 7 *Let $m \geq 3$ and $|N|$ be even. There exists no rule satisfying preference selection, anonymity and K -SP.*

Theorem 8 *Let $m \geq 4$ and $|N|$ be a multiple of 3. There exists no rule satisfying preference selection, anonymity and K -SP.*

Note that the above two theorems exclude situations where there are both three alternatives and three agents. In this case, it turns out that preference selection and K -SP together with either neutrality or anonymity characterize families of rules which are reminiscent to Black's median rules. In other words, then it is possible to choose a representative agent while assuring K -SP and either neutrality or anonymity.

For three alternatives and three agents we denote by \triangle the triangular profiles where agents' preferences are K -equidistant from each other (e.g. (abc, cab, bca)), as illustrated in Figure 5. Triangular profiles correspond exactly to the ones where no Condorcet winner exists, i.e. there is no $x \in A$ such that for any $y \in A \setminus \{x\}$ at least two agents strictly prefer x to y . Note that there are 12 triangular profiles and neutrality divides them into two sets of 6

profiles, i.e. $\Delta = \Delta' + \Delta''$ where Δ' contains (abc, cab, bca) and all profiles obtained from it by permuting alternatives, and where Δ'' contains (acb, bac, cba) and all profiles obtained from it by permuting alternatives. Similarly anonymity divides Δ into two sets of 6 profiles, i.e. $\Delta = \hat{\Delta}' + \hat{\Delta}''$ where $\hat{\Delta}'$ contains (abc, cab, bca) and all profiles obtained from it by permuting agents, and where $\hat{\Delta}''$ contains (acb, bac, cba) and all profiles obtained from it by permuting agents.

Proposition 4 *Let $N = \{1, 2, 3\}$ and $A = \{a, b, c\}$.*

- (i) *A non-dictatorial rule f satisfies preference selection, neutrality and K-SP if and only if there exist $i, j \in N$ (where $i = j$ is possible) such that for all $R_N \in \Delta'$, $f(R_N) = R_i$, for all $R_N \in \Delta''$, $f(R_N) = R_j$, and for all $R_N \in \mathcal{R}^N \setminus \Delta$, the median is chosen, i.e. $f(R_N) = R_i$ if $R_i \in [R_j, R_k]$ (where $N = \{i, j, k\}$) and $[R_j, R_k]$ is the unique shortest path from R_j to R_k to which R_i belongs to. In this case, we say that f is median rule with agent-based tie-breaking.*
- (ii) *A rule f satisfies preference selection, anonymity and K-SP if and only if there exist $\hat{R}'_0 \in \{abc, cab, bca\}$ and $\hat{R}''_0 \in \{acb, bac, cba\}$ such that for all $R_N \in \hat{\Delta}'$, $f(R_N) = \hat{R}'_0$, for all $R_N \in \hat{\Delta}''$, $f(R_N) = \hat{R}''_0$, and for all $R_N \in \mathcal{R}^N \setminus \Delta$, the median is chosen, i.e. $f(R_N) = R_i$ if $R_i \in [R_j, R_k]$ (where $N = \{i, j, k\}$) and $[R_j, R_k]$ is the unique shortest path from R_j to R_k to which R_i belongs to. In this case, we refer to f as a median rule with preference-based tie-breaking.*

Remark 2 *Proposition 4 is a possibility result for three agents and three alternatives where we keep either neutrality or anonymity and always a strict preference order is chosen. Horan et al. [29] and Barberà et al. [11] keep both neutrality and anonymity and obtain only conclusive results whenever Condorcet winners exist. When applied to triangular profiles, the correspondence of Horan et al. [29] selects all three alternatives. Moreover, Horan et al. [29] obtain an impossibility in their Theorem 3 when there are either (i) more than three agents and at least three alternatives or (ii) at least three agents and more than three alternatives. Barberà et al. [11] consider only single-valued choice functions and remain silent for profiles with no Condorcet winners. Now, in Proposition 4, for any non-triangular profile a Condorcet winner exists and the chosen preference order puts this alternative first, and second the Condorcet winner among the two remaining alternatives, which corresponds to the “Condorcet rationalization” in the Appendix of Barberà et al. [11].*

In contrast to these two contributions we obtained possibilities for neutrality for any number of agents and any number of alternatives (Theorem 5), and for anonymity for any number of agents and three alternatives (Proposition 3).

Remark 3 *A more general possibility result is obtained in the vein of Black [15] by restricting both agents' and society's preference orders to be single-peaked. More precisely, let $A = \{a_1, \dots, a_m\}$ be ordered according to their indices, i.e. $a_1 < a_2 < \dots < a_m$. Suppose agent preferences are restricted to be single-peaked according to the order $<$, i.e. for any R_i , for its most preferred alternative $t(R_i) \in A$ (the “peak” of R_i), we have for all $x, y \in A$, $x < y < t(R_i)$ or $x > y > t(R_i)$ implies $t(R_i)P_i y P_i x$. For an odd number of agents, there exists a median of the peaks, denoted by $m(R)$, meaning that more than half of the agents' peaks are smaller than or equal to $m(R)$ and more than half of the agents' peaks are greater than or equal to $m(R)$. The society's preference is restricted to be single-peaked with peak $m(R)$. In addition, preferences over alternative pairs with one left of $m(R)$ and one right of $m(R)$ are determined according to majority. Median rules satisfy anonymity, strong unanimity and K-SP. We refer to Ehlers and Storcken [25] for a detailed analysis of the class of Arrovian aggregation rules in single-peaked environments.*

7 Conclusion

We study the problem of aggregating individual preferences into one preference representing society. Applications range from electing an individual to represent the society to choosing the representative consumer of the economy. In such situations the collective preference must possess the same properties as individual preferences and agents would like it to be “as similar as possible” to their own preference.

One main motivation for our work comes from macroeconomics where individuals' preferences (or the society's preference) have a representative consumer (who has preferences as an individual). Thus, if individuals' preferences are strict, they should be aggregated into a strict preference. Allowing indifferences on both sides (individual and society) is important but beyond the scope of this paper and shall be studied in subsequent research. Similarly, restricting both individual and society's preferences may result in possibilities, which we included in a remark for single-peaked environments at the end of the section on preference selection, but similarly this requires significant more research.

In such settings fairness is important. We explored the tradeoff between anonymity (where agents are treated equally) and neutrality (where alternatives are treated equally), two fundamental properties which are generally impossible to jointly satisfy. As basic requirements we consider strong unanimity, an analogue of Pareto efficiency, and Kemeny strategy-proofness (K-SP), whereby any agent prefers preferences which are closer to his own in terms of Kemeny distance. Dispensing with anonymity and keeping neutrality, we proposed semi-dictator rules and showed they are the first non-trivial family to satisfy strong unanimity, neutrality

and K-SP. Furthermore, for two agents these properties characterize semi-dictator rules. Dispensing with neutrality and keeping anonymity, we found a computer-aided solution to the existence of a rule satisfying strong unanimity, anonymity and K-SP when there are two agents and four alternatives. For three alternatives and an arbitrary number of agents, we provided two families of rules satisfying the desired properties. Finally, for three alternatives and three agents, we characterized median rules with tie-breaking via preference selection, K-SP and either neutrality or anonymity.

When choosing the society’s preference, we must decide how to resolve the anonymity-neutrality fairness tradeoff. If the society finds equal treatment of agents to be more important than equal treatment of alternatives, then neutrality should be dropped and anonymity maintained; otherwise, the opposite should occur. Our results help to clarify the consequences of such a judgment call as regards the design of efficient and strategy-proof aggregation rules.

An interesting avenue of future research would explore how various relaxations of neutrality affect the results of the paper. Consistent with Bartholdi et al.’s [18] and Kivinen’s [32] treatment of anonymity, we could restrict the neutrality criterion to a subset of alternative permutations. For example, we could substitute the full neutrality criterion with a weaker, “local” version that focuses on pairs of alternatives. This form of local neutrality would only require that, for any pair of alternatives, if the identities of two alternatives are flipped then their identities in the social orderings should also be flipped. Relaxing neutrality in this way holds normative appeal and might lead to new possibilities.

Appendix

Below we provide all proofs of our results in the main text.

Proof of Theorem 1: The proof is a straightforward adaptation of the argument of Theorem 1 in Moulin [40].

First we prove necessity. Suppose that $n = k \cdot p$ for some integers k and p such that $p \leq m$. We will show that no anonymous and neutral rule exists. Consider the profile R_N

satisfying (where bold fonts are added for clarity):

$$\begin{aligned}
R_1 = R_{p+1} = R_{2p+1} = \dots R_{(k-1)p+1} &= \mathbf{a_1 a_2 \dots a_p} a_{p+1} \dots a_m \\
R_2 = R_{p+2} = R_{2p+2} = \dots R_{(k-1)p+2} &= \mathbf{a_2 \dots a_{p-1} a_1} a_{p+1} \dots a_m \\
R_3 = R_{p+3} = R_{2p+3} = \dots R_{(k-1)p+3} &= \mathbf{a_3 \dots a_p a_1 a_2} a_{p+1} \dots a_m \\
&\vdots \\
R_p = R_{p+p} = R_{2p+p} = \dots R_{(k-1)p+p} &= \mathbf{a_p a_1 \dots a_{p-1}} a_{p+1} \dots a_m.
\end{aligned}$$

Now define the permutation $\pi : A \mapsto A$ as follows:²³

$$\pi(k) = \begin{cases} \text{mod}_p(k+1) & \text{if } k \in \{1, 2, \dots, p\} \\ k & \text{otherwise.} \end{cases}$$

Simple algebra yields $f(\pi R_N) = f(R_{\sigma(N)})$, where the permutation $\sigma : N \mapsto N$ is given by: for all $l \in \{0, \dots, k-1\}$ and all $i \in \{1, \dots, p\}$,

$$\sigma(lp + i) = lp + \text{mod}_p(i+1).$$

Anonymity requires $f(R_{\sigma(N)}) = f(R_N)$ whereas neutrality requires $f(\pi R_N) = \pi f(R_N) \neq f(R_N)$. This contradicts $f(\pi R_N) = f(R_{\sigma(N)})$.

We now prove sufficiency. Suppose every prime factor of n is greater than m . This means that it is not possible to write $n = k \cdot p$ for some integers k, p such that $p \leq m$. We proceed by displaying a rule that is anonymous and neutral. Given a profile R_N and $a \in A$, define the quantity

$$l(R_N, a) = |\{i \in N : a \text{ ranked last by agent } i \text{ in profile } R_N\}|,$$

i.e., the number of agents who rank a last in profile R_N . In addition, given a profile R_N and $B \subseteq A$, define the

$$L(R_N, B) = \{a \in B : a = \arg \max_{b \in B} l(R_N|_B, b)\},$$

i.e., the set of alternatives attaining the maximum of function $l(R_N|_B, \cdot)$ over set B .

Suppose there exists $B^* \subseteq A$ with $|B^*| > 1$ with the property that, for all $a \in B^*$, there exists the same number k^* of agents ranking a last in R_N . This implies that $n = k^* \cdot |B^*|$, which, since $|B^*| \leq m$, contradicts the stated hypothesis on n and m . Hence, for all $B \subseteq A$,

²³Here $\text{mod}_p(k+1)$ denotes the number modular to p , i.e. $\text{mod}_p(p+1) = 1$ and $\text{mod}_p(k+1) = k+1$ if $k < p$.

we have:

$$L(R_N, B) \subseteq B \text{ and } \{L(R_N, B) = B \Leftrightarrow |B| = 1\}.$$

As a result, given any $B \subseteq A$, the decreasing sequence

$$\begin{aligned} B_0 &= B, \\ B_t &= B_{t-1} \setminus L(R_N, B_{t-1}), \quad t = 1, 2, \dots \end{aligned}$$

will converge to a singleton for some $t \in \{1, 2, \dots, m\}$. We call this alternative $a^*(B)$.

Given a rule f and a profile R_N , let $f_k(R_N)$ denote the k th ranked alternative in $f(R_N)$. Now, define the aggregation rule f^* as the output of the following algorithm:

Input: R_N

1. Initialize $A_0 = A$.
2. For $k = 1, 2, \dots, m$
 - (a) Set $f_k^*(R_N) = a^*(A_{k-1}) \equiv a_k$.
 - (b) Set $A_k = A_{k-1} \setminus a_k$.

Output: $f^*(R_N) = a_1 a_2 \dots a_m$

The above algorithm is well-defined and terminates at $k = m$, since at every k the alternative $a^*(A_{k-1})$ is well-defined. The rule f^* is anonymous and neutral. \square

Proof of Proposition 2: Suppose rule f is locally K-SP and Min-SP but not K-SP. Then there exists an agent i and profile $R_N = (R_i, R_{-i})$ such that $\delta(R_i, f(R'_i, R_{-i})) < \delta(R_i, f(R_N))$, for some R'_i satisfying $\delta(R_i, R'_i) > 1$.

Denote $f(R'_i, R_{-i}) = R^*$. By Min-SP, $f(R^*, R_{-i}) = R^*$. Suppose $\delta(R_i, R^*) = T - 1$ and consider a shortest path between R_i and R^* , which we denote $\{R_i = R^1, R^2, \dots, R^T = R^*\}$. To avoid cumbersome notation, let $x^t \equiv f(R^t, R_{-i})$ for all $t = 1, 2, \dots, T$. By assumption, we have $x^1 = f(R_N)$ and $x^T = R^*$.

We will show by backwards induction that $\delta(R^t, x^t) \leq \delta(R^t, R^*)$ for all t . The induction basis $t = T$ follows trivially because $x^T = R^*$. Suppose $\delta(R^k, x^k) \leq \delta(R^k, R^*)$ for all $k = t, t + 1, \dots, T$. By local K-SP at (R^{t-1}, R_{-i}) and the induction hypothesis applied to $k = t$:

$$\delta(R^{t-1}, x^{t-1}) \leq \delta(R^{t-1}, x^t) \leq 1 + \delta(R^t, x^t) \leq 1 + \delta(R^t, R^*) = \delta(R^{t-1}, R^*).$$

Thus, the induction step is complete, implying that $\delta(R^t, x^t) \leq \delta(R^t, R^*)$ for all $t = 1, 2, \dots, T$. When applied to $t = 1$ this yields $\delta(R_i, f(R_N)) \leq \delta(R_i, R^*)$, which is a contradiction to $f(R'_i, R_{-i}) = R^*$ and $\delta(R_i, f(R'_i, R_{-i})) < \delta(R_i, f(R_N))$. \square

Proof of Theorem 4:

Note that the (if) direction is a special case of Theorem 5 which we show later.

The proof of the (only if) direction proceeds in three steps.

1. First, we prove the characterization when $m = 3$.
2. Then, we use Step 1 to prove the characterization for $m = 4$.
3. Using Step 2 as a base case, we prove the characterization for $m \geq 4$ by induction.

Step 1: The case $m = 3$.

Suppose $N = \{1, 2\}$ and $A = \{a, b, c\}$.

For the (only if) direction, let f satisfy the properties. Focusing on agent 1, we consider the following three rules.

- (a) For all $R_N = (R_1, R_2) \in \mathcal{R}^N$, $f^0(R_N) = R_1$; or
- (b) For all R_N with $R_1 = a_1 a_2 a_3$, $f^1(R_N) = (a_1 R_2|_{\{a_2, a_3\}})$ (where $R_2|_{\{a_2, a_3\}}$ denotes the restriction of R_2 to a_2 and a_3); or
- (c) For all R_N with $R_1 = a_1 a_2 a_3$, $f^2(R_N) = (R_2|_{\{a_1, a_2\}} a_3)$ (where $R_2|_{\{a_1, a_2\}}$ denotes the restriction of R_2 to a_1 and a_2).

Denote the corresponding rules where agent 2 plays the role of agent 1 and vice versa by g^0 , g^1 and g^2 .

Recall that a profile R_N is opposite if $R_2 = -R_1$. Consider $f(abc, cba)$, and without loss of generality, let $f(abc, cba) \in \{acb, abc, bac\}$. If this is not the case, then focus on agent 2 and rules g^0, g^1, g^2 and apply similar reasoning. We distinguish between three cases.

1. $f(abc, cba) = abc$. Then by neutrality, for any opposite profile R_N , agent 1's preference is chosen, i.e. $f(R, -R) = R = f^0(R_N)$ for all $R \in \mathcal{R}$. Since $f(R_1, -R_1) = R_1$ for any choice of R_1 , K-SP applied to agent 2 implies that, i.e., $f(R_1, R_2) = R_1 = f^0(R_1, R_2)$ for all $R_1, R_2 \in \mathcal{R}$.
2. $f(abc, cba) = acb$. Then by neutrality, for any opposite profile R_N we have $f(R_N) = f^1(R_N)$. If $R_N = (R_1, R_2)$ is not opposite, then consider $f(R_1, -R_1) = f^1(R_1, -R_1)$ and $f(R_2, -R_2) = f^1(R_2, -R_2)$. Now if $R_2 \neq -R_1$ is on the half circle that links R_1

to $-R_1$ which includes $f^1(R_1, -R_1)$, then by strong unanimity we have $f(R_1, R_2) \in \{R_1, f^1(R_1, -R_1), R_2\}$. K-SP applied to agent 2 at profile $(R_1, -R_1)$ yields $f(R_1, R_2) \neq R_2$. Similarly, K-SP applied to agent 2 at profile (R_1, R_2) implies $f(R_1, R_2) \neq R_1$. Hence we conclude $f(R_1, R_2) = f^1(R_1, R_2)$.

Otherwise, $R_2 \neq -R_1$ is not on the half circle containing $f^1(R_1, -R_1)$. For clarity, and without loss of generality (due to neutrality) suppose $R_1 = abc$, so that $f(abc, cba) = acb$ and $R_2 \in \{bac, bca\}$. If $R_2 = bca$, then by neutrality, $f(-R_2, R_2) = abc$, and by K-SP, $f(R_N) = abc = f^1(R_N)$, the desired conclusion. If $R_2 = bac$, then from the previous fact, $f(acb, bca) = abc$. By K-SP and strong unanimity, $f(acb, R_2) = abc$, and using again K-SP and strong unanimity, we obtain $f(R_N) = abc = f^1(R_N)$, the desired conclusion.

3. $f(abc, cba) = bac$. Note that in this case $f(abc, cba) = bac = f^2(abc, cba)$. Then using similar arguments as in Case 2 it follows that $f(R_N) = f^2(R_N)$ for all R_N .

□

Step 2: The case $m = 4$.

Let $A = \{a, b, c, d\}$. For the (only if) direction, let f satisfy the properties. Let \underline{f} denote the rule where f is restricted to the domain where both agents rank at the bottom the same alternative. By strong unanimity, also \underline{f} ranks at the bottom this alternative. Thus, \underline{f} is a two agents-three alternatives rule. Furthermore, neutrality implies that the same type of rule is chosen when the two agents rank the same alternative at the bottom. Similarly we denote by \overline{f} the rule where f is restricted to the domain where both agents rank at the top the same alternative. We consider three cases: (I) $\underline{f} = f^1$, (II) $\underline{f} = f^0$ and (III) $\underline{f} = f^2$ (as $\underline{f} \in \{g^0, g^1, g^2\}$ is analogous to the one by switching the roles of agent 1 and agent 2).

(I) $\underline{f} = f^1$:

Suppose that \underline{f} is of type f^1 . We show that f must be a semi-dictator rule with semi dictator 1 and agent 2 chooses the preference in f of the second and third alternatives of 1's preference.

First, we show that $f_1(R_N) = \text{top}(R_1)$ for all R_N . Suppose not, i.e. $f_1(R_N) \neq \text{top}(R_1)$. Then by K-SP, $f(R_1, f(R_N)) = f(R_N)$. Let $R'_1 : \text{top}(R_1)f(R_N)|_{A \setminus \{\text{top}(R_1)\}}$. Then $R'_1 \in [R_1, f(R_N)]$ and by K-SP, $f(R'_1, f(R_N)) = f(R_N)$. If R'_1 and $f(R_N)$ rank the same alternative at the bottom, then this is a contradiction to $\underline{f}(R'_1, f(R_N)) = f^1(R'_1, f(R_N))$ and

$$f_1(R'_1, f(R_N)) = f_1(R_N) \neq \text{top}(R_1) = \text{top}(R'_1).$$

Thus, $\text{bot}(f(R_N)) = \text{top}(R_1)$. Let $R'_2 : f_1(R_N)f_2(R_N)f_4(R_N)f_3(R_N)$. Note that $f(R'_1, f(R_N))$ is of K-distance one to R'_2 . Thus, $f(R'_1, R'_2)$ is of distance at most one to R'_2 , which implies $f_1(R'_1, R'_2) \neq \text{top}(R'_1)$, a contradiction because $f_4(R'_1, f(R_N)) = f_4(R_N) = \text{top}(R_1) = \text{top}(R'_1)$, $\text{bot}(R'_1) = \text{bot}(R'_2)$, $f(R'_1, R'_2) = \underline{f}(R'_1, R'_2)$ and $\underline{f} = f^1$.

Second, we show that $f_2(R_N) \neq \text{bot}(R_1)$ for all R_N . Suppose that $f_2(R_N) = \text{bot}(R_1) \neq \text{bot}(R_2)$ (as otherwise we have a contradiction to the fact that \underline{f} is of type f^1). Let $R_1 : abcd$. By K-SP, $f(R_1, f(R_N)) = f(R_N)$ and $f(R_N) \in \{adbc, adcb\}$. Thus, without loss of generality, we may suppose $R_2 = f(R_N)$.

Case 1: $f(R_N) : adcb$.

Then $\delta(R_1, f(R_N)) = 3$. Let $R'_1 : bacd$. Then from $\text{top}(f(R'_1, f(R_N))) = \text{top}(R'_1) = b$ and by strong unanimity, $f(R'_1, f(R_N)) \in \{badc, bacd\}$. But then $\delta(R_1, f(R'_1, f(R_N))) \leq 2$, a contradiction to K-SP.

Case 2: $f(R_N) : adbc$.

Let $\hat{R}_1 : acbd$ and $\hat{R}_2 : adcb$. Note that $\hat{R}_N = R_N^{b \leftrightarrow c}$ and by neutrality, $f(\hat{R}_N) = adcb$. Consider $(\hat{R}_1, f(R_N))$. Then $\delta(f(R_N), f(\hat{R}_N)) = 1$ and by K-SP (as agent 2 could deviate from $(\hat{R}_1, f(R_N))$ to $\hat{R}_N = (\hat{R}_1, \hat{R}_2)$), $\delta(f(R_N), f(\hat{R}_1, f(R_N))) \leq 1$. Thus, $f(\hat{R}_1, f(R_N)) \in \{adbc, dabc, abdc, adcb\}$. Then

- (i) $f(\hat{R}_1, f(R_N)) = adbc$ implies that $\bar{f} = g^0$ and $f(R_1, adcb) = adcb$, which yields a contradiction as in Case 1;
- (ii) $f(\hat{R}_1, f(R_N)) = dabc$ contradicts the fact $f_1(\hat{R}_1, f(R_N)) = \text{top}(\hat{R}_1) = a$;
- (iii) $f(\hat{R}_1, f(R_N)) = abdc$ implies that \bar{f} is of type g^2 which implies for $\bar{R}_1 : abcd$ and $\bar{R}_2 : acbd$ we have both $f(\bar{R}_N) = \bar{f}(\bar{R}_N) = g^2(\bar{R}_N) = abcd$ and $f(\bar{R}_N) = \underline{f}(\bar{R}_N) = f^1(\bar{R}_N) = acbd$, a contradiction; and
- (iv) $f(\hat{R}_1, f(R_N)) = adcb$ implies that \bar{f} is of type g^1 .

For (iv) we derive a contradiction in three steps. In the first step we show that $f_2(R_N) = \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}})$ for any profile R_N . In the second step then we show that either agent 1 always chooses the third alternative in $f(R_N)$ or agent 2 always chooses the third alternative. In the third step we show that f violates K-SP (and therefore, (iv) cannot occur).

In the first step we show that $f_2(R_N) = \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}})$ for any profile R_N . Suppose $f_2(R_N) \neq \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}})$. By K-SP and neutrality, without loss of generality, we may suppose $R_1 = f(R_N) = abcd$. By $\bar{f} = g^1$, we have $a = \text{top}(R_1) \neq \text{top}(R_2)$ and $\text{top}(R_2) = \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}}) \neq a, b$ (as $b = f_2(R_N) \neq \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}})$). Similarly, by $\underline{f} = f^1$, we must have $\text{bot}(R_2) \neq d = \text{bot}(R_1)$. We distinguish two subcases ($\text{top}(R_2) = c$ or $\text{top}(R_2) = d$):

if $\text{top}(R_2) = c$, then for $R'_2 : cabd$ we have $R'_2 \in [R_2, f(R_N)]$ (as $R_1 = f(R_N) = abcd$) and $f(R_1, R'_2) = f(R_N) = abcd$ which is a contradiction as $\underline{f} = f^1$ and $f(R_1, R'_2) = \underline{f}(R_1, R'_2) = acbd$; and if $\text{top}(R_2) = d$, then for $R'_2 : dabc$ we have $R'_2 \in [R_2, f(R_N)]$, $f(R_1, R'_2) = f(R_N) = abcd$ and $\delta(R'_2, f(R_N)) = 3$ which yields a contradiction to K-SP as for $R'_2 : adbc$ we have (from $\bar{f} = g^1$) $f(R_1, R'_2) = \bar{f}(R_1, R'_2) = adbc$ and $\delta(R'_2, f(R_1, R'_2)) = 1$.

For the second step, consider $R'_1 : abcd$, $R'_2 : abdc$ and $R'_N = (R'_1, R'_2)$. By strong unanimity, $f(R'_N) = R'_1$ or $f(R'_N) = R'_2$. We show that if $f(R'_N) = R'_1$, then agent 1 always chooses the third alternative, and if $f(R'_N) = R'_2$, then agent 2 always chooses the third alternative. Without loss of generality, let $f(R'_N) = R'_1$. Let R_N be an arbitrary profile. By the above, we have $f_1(R_N) = \text{top}(R_1)$ and $f_2(R_N) = \text{top}(R_2|_{A \setminus \{\text{top}(R_1)\}})$. Suppose $f_3(R_N) \neq \text{top}(R_1|_{A \setminus \{f_1(R_N), f_2(R_N)\}})$. Then by K-SP, $f(R_1, f(R_N)) = f(R_N)$. Let $R'_1 = f_1(R_N)f_2(R_N)f_4(R_N)f_3(R_N)$. Now by neutrality and our assumption, $f(R'_1, f(R_N)) = R'_1$. Note that $f_4(R_N) = \text{top}(R_1|_{A \setminus \{f_1(R_N), f_2(R_N)\}})$, and $\delta(R_1, R'_1) < \delta(R_1, f(R_N))$, which is a contradiction to K-SP.

In the third step we show that f violates K-SP. Let $R_1 : abcd$ and $R_2 : adcb$. If $f(R'_N) = R'_1$, then by the above, $f(R_N) : adbc$. Let $\hat{R}_1 : bacd$. Then $f(\hat{R}_1, R_2) = bacd$, $\delta(R_1, f(\hat{R}_1, R_2)) = 1 < \delta(R_1, f(R_N))$, which is a contradiction to K-SP. If $f(R'_N) = R'_2$, then as above it can be shown agent 2 always chooses the third alternative in $f(R_N)$. But then consider $R_1 : abcd$ and $R_2 : adcb$. Then $f(R_N) = adcb$ but for $\hat{R}_1 : bacd$ we have $f(\hat{R}_1, R_2) : badc$, $\delta(R_1, f(\hat{R}_1, R_2)) = 2 < 3 = \delta(R_1, f(R_N))$. Thus, (iv) also leads to a contradiction.

We have shown $f_1(R_N) = \text{top}(R_1)$ and $f_2(R_N) \neq \text{bot}(R_1)$ for all R_N . Now if $\text{bot}(R_2) = \text{bot}(R_1)$, then we have $f(R_N) = \underline{f}(R_N)$ where $f_1(R_N) = \text{top}(R_1)$ and $f_4(R_N) = \text{bot}(R_1)$, and by $\underline{f} = f^1$, R_2 decides the ranking over the second and third alternative in R_1 (which is the desired conclusion).

Before treating the remaining case $\text{bot}(R_2) \neq \text{bot}(R_1)$, as an intermediary step, we show $\bar{f} = f^2$. Consider $R_1 : abcd$ and $R_2 : acbd$. Then $f(R_N) = \underline{f}(R_N) = acbd$. As also $f(R_N) = \bar{f}(R_N)$, we obtain $\bar{f} \neq f^0, f^1, g^2$. Thus, $\bar{f} \in \{f^2, g^0, g^1\}$. If $\bar{f} \in \{g^0, g^1\}$, then for $R_1 : abcd$ and $R_2 : adcb$ we have $f_2(R_N) = d$, a contradiction to $f_2(R_N) \neq \text{bot}(R_1)$. Hence, we obtain $\bar{f} = f^2$.

Finally, suppose $\text{bot}(R_1) \neq \text{bot}(R_2)$. We show $f_4(R_N) = \text{bot}(R_1)$. Suppose $f_4(R_N) \neq \text{bot}(R_1)$ and by neutrality, without loss of generality, let $R_1 = abcd$. Then $f_1(R_N) = a$ and by $f_2(R_N) \neq \text{bot}(R_1)$, $f_3(R_N) = d$. Thus, $f(R_N) \in \{abdc, acdb\}$.

If $f(R_N) = acdb$, then by K-SP, $f(R_1, f(R_N)) = acdb \neq \bar{f}(R_1, f(R_N)) = f^2(R_1, f(R_N))$, which is a contradiction to the above.

Hence, $f(R_N) = abdc$. Then by cP_1d and strong unanimity, dP_2c . As $\text{top}(R_2) \neq a$, we have

then $\text{top}(R_2) \in \{b, d\}$. If $\text{top}(R_2) = d$, then dP_2b . By K-SP, $f(f(R_N), R_2) = f(R_N) = abdc$. But now for $R'_2 : adbc$ we have (from $\underline{f} = f^1$) $f(f(R_N), R'_2) = \underline{f}(f(R_N), R'_2) = adbc$ which is a contradiction to K-SP as

$$\delta(f(R_2, f(f(R_N), R'_2))) = \delta(R_2, adbc) < \delta(R_2, abdc) = \delta(R_2, f(R_N)) = \delta(R_2, f(f(R_N), R_2)),$$

where the inequality follows from dP_2b .

If $\text{top}(R_2) = b$, then from strong unanimity, $f(R_N) = abdc$ and cP_1d we obtain dP_2c . Now for $R'_2 : bdca$ we have $f_1(R_N) = a = f_1(R_1, R'_2)$ and by K-SP, $f(R_1, R'_2) = f(R_N) = abdc$ (where agent 2 is misreporting). But then by K-SP, $f(R_1, abdc) = abdc \neq abcd = f^2(R_1, abdc) = \bar{f}(R_1, abdc)$, a contradiction.

Hence, we have shown that for any profile R_N , $f_1(R_N) = \text{top}(R_1)$ and $f_4(R_N) = \text{bot}(R_1)$. As $\underline{f} = f^1$ it follows that agent 2 chooses the preference in $f(R_N)$ of the second and third alternatives of R_1 (as by K-SP we may suppose $\text{top}(R_2) = f_1(R_N)$ and $\text{bot}(R_2) = f_4(R_N)$).

(II) $\underline{f} = f^0$.

We show that for any profile R_N , $f(R_N)$ restricted to its first three alternatives coincides with R_1 restricted to these alternatives. Again by K-SP, without loss of generality, let $R_2 = f(R_N)$. If $f(R_N)|_{A \setminus \{f_4(R_N)\}} \neq R_1|_{A \setminus \{f_4(R_N)\}}$, then let $R'_1 : R_1|_{A \setminus \{f_4(R_N)\}} f_4(R_N)$ and then $f(R'_1, R_2) = \underline{f}(R'_1, R_2) = f^0(R'_1, R_2) = R'_1$ which is a contradiction to K-SP as by $f_4(R_N) = f_4(R'_1, R_2)$ we have $\delta(R_1, f(R'_1, R_2)) < \delta(R_1, f(R_N))$.

Hence, for any R_N ,

$$f(R_N) : R_1|_{A \setminus \{f_4(R_N)\}} f_4(R_N). \quad (9)$$

Next we show $f_1(R_N) = \text{top}(R_1)$. If $f_1(R_N) \neq \text{top}(R_1)$, then by the previous fact, $f_4(R_N) = \text{top}(R_1)$ and $f_3(R_N) = \text{bot}(R_1)$. Let $R'_2 : f_1(R_N) f_2(R_N) f_4(R_N) f_3(R_N)$. As $\delta(R'_2, f(R_N)) = 1$, $f(R_1, R'_2)$ must be of K-distance one or zero to R'_2 which implies $\text{top}(R_1) \neq f_1(R_1, R'_2)$ and by (9), $f_4(R_1, R'_2) = \text{top}(R_1)$ and $f(R_1, R'_2) = f(R_N)$. Then $\delta(R_1, f(R_1, R'_2)) = 3$. Let $R'_1 : R_1|_{A \setminus \{f_3(R_N)\}} f_3(R_N)$. Then $f(R'_1, R'_2) = \underline{f}(R'_1, R'_2) = f^0(R'_1, R'_2) = R'_1$ and $\delta(R_1, f(R'_1, R'_2)) < 3 = \delta(R_1, f(R_N)) = \delta(R_1, f(R_1, R'_2))$, a contradiction to K-SP. We further show $f_2(R_N) = \text{top}(R_1|_{A \setminus \{\text{top}(R_1)\}})$. If $f_2(R_N) \neq \text{top}(R_1|_{A \setminus \{\text{top}(R_1)\}})$, then for $R_1 = abcd$ we obtain from $f_1(R_N) = \text{top}(R_1)$ and (9) that $f(R_N) = acdb$. Again let $R_2 = f(R_N)$. Then $\bar{f} \in \{g^0, g^2\}$. By considering the profile $\hat{R}_N = (abcd, acbd)$ we then get a contradiction to $\bar{f} = g^0$ as $f(\hat{R}_N) = \underline{f}(\hat{R}_N) = f^0(\hat{R}_N) = \hat{R}_1$ and $f(\hat{R}_N) = \bar{f}(\hat{R}_N) = g^0(\hat{R}_N) = \hat{R}_2$. Hence, $\bar{f} = g^2$. Recall that $f(R_N) = acdb = R_2$ and $\delta(R_1, f(R_N)) = 2$. Let $R'_1 : bacd$. But then $f_1(R'_1, R_2) = \text{top}(R'_1) = b$. By strong unanimity then $f(R'_1, R_2) = bacd$ and $\delta(R_1, f(R'_1, R_2)) = 1$, a contradiction to K-SP.

We have shown that for any profile R_N , $f_1(R_N) = \text{top}(R_1)$ and $f_2(R_N) = \text{top}(R_1|_{A \setminus \{\text{top}(R_1)\}})$. Now considering $R'_1 = abcd$ and $R'_2 = abdc$ we have $f(R'_N) \in \{abcd, abdc\}$. If $f(R'_N) = abcd$, then R_1 always decides the ranking of the third and fourth alternative in $f(R_N)$: for any profile R_N by neutrality we may suppose $R_1 = abcd$; then $f_1(R_N) = a$ and $f_2(R_N) = b$; again by K-SP we may choose $R_2 = f(R_N)$ and obtain $f(R_N) = f(R_1, f(R_N))$; hence, $(R_1, f(R_N)) = R'$ and we must have $f(R_N) = abcd = R_1$, the desired conclusion. If $f(R'_N) = abdc$, then similarly it can be shown that R_2 always decides the ranking of the third and fourth alternative in $f(R_N)$.

(III) $\underline{f} = f^2$.

We show that for any profile R_N , $\{f_3(R_N), f_4(R_N)\}$ consists of the third and fourth ranked alternative in R_1 and R_2 decides the ranking of the first two alternatives in $f(R_N)$.

First, we show $f_1(R_N) \neq \text{bot}(R_1)$. Let $R_1 : abcd$ and suppose $f_1(R_N) = d$ and $R_2 = f(R_N)$ (by K-SP). By $\underline{f} = f^2$, we must have $\text{bot}(R_2) \neq d$. Let $R'_1 : R_1|_{A \setminus \{f_4(R_N)\}} f_4(R_N)$. Then $f(R'_1, R_2) = \underline{f}(R'_1, R_2)$ which implies $f_3(R'_1, R_2) = d$ and $f_4(R'_1, R_2) = f_4(R_N)$. But then by $f_1(R_N) = d$, $\delta(R_1, f(R_N)) > \delta(R_1, f(R'_1, R_2))$, a contradiction to K-SP.

Second, we show $f_2(R_N) \neq \text{bot}(R_1)$. Let $R_1 : abcd$ and suppose $f_2(R_N) = d$ and $R_2 = f(R_N)$ (by K-SP). By $\underline{f} = f^2$, we must have $\text{bot}(R_2) \neq d$. Let $R'_1 : R_1|_{A \setminus \{f_4(R_N)\}} f_4(R_N)$. Then $f(R'_1, R_2) = \underline{f}(R'_1, R_2)$ which implies $f_3(R'_1, R_2) = d$ and $f_4(R'_1, R_2) = f_4(R_N)$. But then by $f_1(R_N) = d$, $f(R_N) = R_2$ and thus, $f(R_N)|_{A \setminus \{d, f_4(R_N)\}} = f(R'_1, R_2)|_{A \setminus \{d, f_4(R_N)\}}$, we obtain $\delta(R_1, f(R_N)) > \delta(R_1, f(R'_1, R_2))$, a contradiction to K-SP.

Thus, we have shown $\text{bot}(R_1) \in \{f_3(R_N), f_4(R_N)\}$. We show that the third alternative in R_1 cannot be ranked first or second in $f(R_N)$, i.e. for $R_1 : abcd$ we have $\{c, d\} = \{f_3(R_N), f_4(R_N)\}$ (as $\text{bot}(R_1) \neq f_1(R_N), f_2(R_N)$). Again, by K-SP, without loss of generality, $R_2 = f(R_N)$. If $\text{bot}(R_2) = d = \text{bot}(R_1)$, then this is obvious from $\underline{f} = f^2$. Thus, let $\text{bot}(R_2) \neq d$. If $\text{bot}(R_2) = c$, then this follows from strong unanimity as $d \in \{f_3(R_N), f_4(R_N)\}$. Thus, let $\text{bot}(R_2) \in \{a, b\}$.

Third, we show $f_1(R_N) \neq c$. If $f_1(R_N) = c$, then from $f_4(R_N) = \text{bot}(R_2) \neq c, d$ and $\text{bot}(R_1) = d \in \{f_3(R_N), f_4(R_N)\}$ we obtain $f(R_N) \in \{cadb, cbda\}$. Let $R'_2 : R_2|_{\{a, b, c\}} d$. Then $f(R_1, R'_2) = \underline{f}(R_1, R'_2) = R_2|_{\{a, b\}} cd$. But then $\delta(R'_2, f(R_N)) = 1 < \delta(R'_2, f(R_1, R'_2))$ (as $\text{top}(R'_2) = c$), a contradiction to K-SP.

Fourth, we show $f_2(R_N) \neq c$ in four steps: in the first step we show $\bar{f} = g^2$; in the second step we show $\text{bot}(R_2) \notin \{f_1(R_N), f_2(R_N)\}$; in the third setp we show that for any profile the third ranked alternative in R_2 can never be chosen first in $f(R_N)$; and in the fourth step we obtain a contradiction by showing that for certain profiles no alternative can be ranked first (using the first three steps).

In showing the first step, $f_2(R_N) = c$ implies $f(R_N) \in \{acdb, bcda\}$ (as $d = \text{bot}(R_1) \neq \text{bot}(R_2) = f_4(R_N)$ and $d \in \{f_1(R_N), f_2(R_N)\}$). If $f(R_N) = acdb$, then $\bar{f} = g^0$ or $\bar{f} = g^2$. For $\bar{f} = g^0$, let $R'_1 = abdc$ and then $f(R'_1, R_2) = \bar{f}(R'_1, R_2) = g^0(R'_1, R_2) = acdb$ which is a contradiction as $c = \text{bot}(R'_1) \notin \{f_1(R'_1, R_2), f_2(R'_1, R_2)\}$. Thus, $f(R_N) = acdb$ implies $\bar{f} = g^2$. We show that $f(R_N) = bcda$ also implies $\bar{f} = g^2$: note that $\delta(R_1, f(R_N)) = 3$ which implies by K-SP for $R'_1 = bacd$ that $f(R'_1, R_2) = bcda$ (as $f_1(R'_1, R_2) = b$, $f(R'_1, R_2)$ must rank c before d by strong unanimity, and d must be ranked before a by K-SP as otherwise agent 1 profitably misreports from $f(R_N) = bcda$); but then $\bar{f} = g^0$ or $\bar{f} = g^2$ and again $\bar{f} = g^0$ yields a contradiction as above. Thus, $\bar{f} = g^2$.

In the second step we show that $\text{bot}(R_2) \notin \{f_1(R_N), f_2(R_N)\}$ for any profile R_N . Again suppose $f_1(R_N) = \text{bot}(R_2)$ and $f(R_N) = R_1 = abcd$. Thus, $\text{bot}(R_2) = a$. For $R'_2 : R_2|_{\{b,c\}} ad$ we have (from $\underline{f} = f^2$) $f(R_1, R'_2) = bacd$ and $\delta(R_2, f(R_1, R'_2)) < \delta(R_2, f(R_N))$, a contradiction to K-SP. Next suppose $f_2(R_N) = \text{bot}(R_2)$ and $f(R_N) = R_1 = abcd$. Thus, $\text{bot}(R_2) = b$. For $R'_2 : aR_2|_{\{c,d\}} b$ we have (from $\underline{f} = g^2$) $f(R_1, R'_2) = acdb$ and $\delta(R_2, f(R_1, R'_2)) < \delta(R_2, f(R_N))$, a contradiction to K-SP.

In the third step we show that the third ranked alternative in R_2 can never be chosen first by $f(R_N)$. Suppose $R_2 = abcd$, $f_1(R_N) = c$ and $R_1 = f(R_N)$ (by K-SP). But then $\text{bot}(R_1) \neq \text{bot}(R_2)$ as otherwise $f(R_N) = \underline{f}(R_N)$ and by $\underline{f} = f^2$, R_2 decides the ranking of the first two alternatives in $f(R_N)$ and $f_1(R_N) \neq c$. Thus, from $f_1(R_N) \neq d$ and $R_1 = f(R_N)$, we have $f(R_N) \in \{cbda, cadb\}$. If $f(R_N) = cbda$, then for $R'_2 = bcda$ we have $f(R_1, R'_2) = \underline{f}(R_1, R'_2) = f^2(R_1, R'_2) = bcda$ and $\delta(R_2, f(R_1, R'_2)) < \delta(R_2, f(R_N))$, a contradiction to K-SP. If $f(R_N) = cadb$, then for $R'_2 = acdb$ we have $f(R_1, R'_2) = \underline{f}(R_1, R'_2) = f^2(R_1, R'_2) = acdb$ and $\delta(R_2, f(R_1, R'_2)) < \delta(R_2, f(R_N))$, a contradiction to K-SP.

In the fourth step we now obtain a contradiction for the profile R_N where $R_1 = abcd$ and $R_2 = dcba$ as by $\text{bot}(R_1) \neq \text{bot}(R_2)$ and $\text{bot}(R_1), \text{bot}(R_2) \in \{f_3(R_N), f_4(R_N)\}$ we have $\{a, d\} = \{f_3(R_N), f_4(R_N)\}$. On the other hand the third ranked alternative in R_1 is c and the third ranked alternative in R_2 is b and $f_1(R_N) \in \{b, c\}$ which is a contradiction as then no alternative can be ranked first in $f(R_N)$ as the third ranked alternative in R_1 and in R_2 are never ranked first by $f(R_N)$. This concludes the proof that $f_2(R_N)$ cannot be the third ranked alternative in R_1 for any profile R_N .

We have shown that $\{f_3(R_N), f_4(R_N)\}$ always consists of the third and fourth ranked alternatives in R_1 . We next show that R_2 determines the ranking of the first two alternatives in $f(R_N)$. Again, by K-SP, without loss of generality, $R_1 = f(R_N)$. If $f(R_N)|_{\{f_1(R_N), f_2(R_N)\}} \neq R_2|_{\{f_1(R_N), f_2(R_N)\}}$, then let $R'_2 : R_2|_{\{f_1(R_N), f_2(R_N)\}} f_3(R_N) f_4(R_N)$ and then $f(R_1, R'_2) = \underline{f}(R_1, R'_2) = f^2(R_1, R'_2) = R'_2$, a contradiction to K-SP. Thus, R_2 always determines the ranking of the first two alternatives. Now considering $R_1 = abcd$ and $R_2 = abdc$ we have $f(R_N) \in \{abcd, abdc\}$.

If $f(R_N) = abcd$, then it can be shown analogously that R_1 always decides the ranking of the third and fourth alternative in $f(R_N)$, and if $f(R_N) = abdc$, then it can be shown analogously that R_2 always decides the ranking of the third and fourth alternative in $f(R_N)$. \square

Note that using the same argument as in Example 2, it follows that agent 2 cannot decide both the ranking of the first and second alternative and the ranking of the third and fourth alternative. This finishes the proof of Theorem 4 for four alternatives.

Step 3: Induction on m .

Now by induction suppose that Theorem 4 is true for $k \geq 4$ alternatives (where $m = k$). Let $A = \{a_1, \dots, a_{k+1}\}$. Let f satisfy the properties. Let \underline{f} denote the rule where f is restricted to the domain where both agents rank at the bottom the same alternative, i.e. $\underline{\mathcal{R}}^N = \{R \in \mathcal{R}^N : \text{bot}(R_1) = \text{bot}(R_2)\}$ and $\underline{f} = f|_{\underline{\mathcal{R}}^N}$. By strong unanimity, for any $R_N \in \underline{\mathcal{R}}^N$, \underline{f} ranks at the bottom the alternative $\text{bot}(R_1) = \text{bot}(R_2)$. Thus, by the induction hypothesis and neutrality, \underline{f} is a semi-dictator rule. Similarly we denote by \bar{f} the rule where f is restricted to the domain where both agents rank at the top the same alternative, and again by the induction hypothesis, \bar{f} is a semi-dictator rule. Note that in a semi-dictator rule with semi-dictator i , for any profile R_N any alternative in R_i can move at most one position up or at most one position down in the chosen ranking.

First, we show that \underline{f} and \bar{f} have the same semi-dictator. Suppose not, say agent 1 is the semi-dictator of \underline{f} and agent 2 is the semi-dictator of \bar{f} . Consider $R_1 : a_1 \dots a_{k+1}$ and $R_2 : a_1 \dots a_{k-3} a_k a_{k-1} a_{k-2} a_{k+1}$. Note that we have $f(R_N) = \underline{f}(R_N) = \bar{f}(R_N)$. Thus, by strong unanimity and the fact that both a_k and a_{k-2} can move at most one position up in $f(R_N)$, we have $f(R_N) \in \{a_1 \dots a_{k-1} a_{k-2} a_k a_{k+1}, a_1 \dots a_{k-1} a_k a_{k-2} a_{k+1}\}$. But then either a_k moves down two positions from R_2 to $f(R_N)$ or a_{k-2} moves two positions down from R_1 to $f(R_N)$, a contradiction.

Hence, \underline{f} and \bar{f} have the same semi-dictator say agent 1. But then the positions p where agent 2 is decisive with $1 < p < k$ coincide in \underline{f} and \bar{f} (by considering profiles R_N where $\text{top}(R_1) = \text{top}(R_2)$ and $\text{bot}(R_1) = \text{bot}(R_2)$ and $f(R_N) = \underline{f}(R_N) = \bar{f}(R_N)$ noting $p+1 \leq k$ by $p < k$). Using the same argument as above, it also follows that if $p = 1$ is a position in \underline{f} where agent 2 is decisive, then position 2 does not belong to \bar{f} .

Now consider any profile R_N . By neutrality, we may suppose $R_1 : a_1 \dots a_{k+1}$. By K-SP again we may suppose $R_2 = f(R_N)$ (because otherwise $R_2 \neq f(R_N)$ and $f(R_N) = f(R_1, f(R_N))$ by K-SP). If $f_{k+1}(R_N) = a_{k+1} = \text{bot}(R_1)$, then $\text{bot}(R_1) = \text{bot}(R_2)$ as $R_2 = f(R_N)$ and $f(R_N) = \underline{f}(R_N)$; and we are done by the induction hypothesis. Otherwise ($f_{k+1}(R_N) \neq a_{k+1}$) we show $f_k(R_N) = a_{k+1}$. If not, i.e. $f_l(R_N) = a_{k+1}$ with $l < k$, then let $R'_1 : f(R_N)|_{A \setminus \{a_{k+1}\}} a_{k+1}$. Then $R'_1 \in [R_1, f(R_N)]$ and by K-SP and strong unanimity,

$f(R'_1, R_2) = f(R_N)$. If $f_1(R_N) = \text{top}(R'_1)$, then this is a contradiction to the fact that agent 1 is the semi-dictator of \bar{f} and a_{k+1} moves more than one position up in $f(R_N)$. If $f_1(R_N) \neq \text{top}(R'_1)$, then $f_1(R_N) = a_{k+1}$ and by $k \geq 4$ we obtain a contradiction to K-SP when we exchange in R'_1 the positions of the last two alternatives, i.e. for $R''_1 : f(R_N)|_{A \setminus \{a_{k+1}, f_{k+1}(R_N)\}} a_{k+1} f_{k+1}(R_N)$ we obtain $f(R''_1, R_2) = \underline{f}(R''_1, R_2)$.

Hence, $f_{k+1}(R_N) \neq a_{k+1}$ implies $f_k(R_N) = a_{k+1}$. Similarly, by considering \bar{f} this also implies $f_{k+1}(R_N) = a_k$ (as otherwise $f_{k+1}(R_N) = a_l$ for $l < k$ and a_l moves more than one position down from R_1 to $f(R_N)$). But then we are done as now we consider $R_2 : a_1 \dots a_{k-1} a_{k+1} a_k$. If $f(R_N) = R_2$, then k is a position in f where agent 2 is decisive and otherwise not. Furthermore, note that the ranking of $f(R_N)$ over $\{a_1, \dots, a_{k-1}\}$ is decided by \underline{f} .

Note that using the same argument as in Example 2, it follows that any two positions where agent 2 decides the ranking must have distance greater than two. \square

Proof of Theorem 5:

Consider a semi-dictator rule $f = f^{(i, P, \mathcal{C})}$. That strong unanimity and neutrality are satisfied is obvious from Definitions 6 and 7, so we focus on proving K-SP. Without loss of generality assume $i = 1$ and $R_1 = a_1 a_2 \dots a_m$. By Proposition 2 it is sufficient to prove that f is Min-SP and Locally K-SP.

As before with strong unanimity, Min-SP follows immediately from Definitions 6 and 7. We turn to proving Local K-SP.

Consider a profile R_N and suppose agent j changes his preferences to R'_j , where $\delta(R_j, R'_j) = 1$. If $j \neq 1$, then by the definition of semi-dictator rules and committees, it is clear that this agent cannot profit from misreporting. In fact, $\delta(R_j, f(R'_j, R_{-j})) - \delta(R_j, f(R_N)) \in \{0, 1\}$.

Thus, suppose $j = 1$, meaning that the semi-dictator is the agent who misreports. Suppose R'_1 is identical to R_1 except that the order of two adjacent alternatives a_k and a_{k+1} is flipped for some $k = 1, 2, \dots, m-1$. We distinguish between four cases:

1. $k-1 \notin P$, $k \notin P$ and $k+1 \notin P$. In this case, $f_l(R_N) = f_l(R'_1, R_{-1})$ for all $l \notin \{k, k+1\}$. Since $k-1, k$ and $k+1$ do not belong to P , it is immediate that $f_k(R_N) = a_k, f_{k+1}(R_N) = a_{k+1}$ and $f_k(R'_1, R_{-1}) = a_{k+1}, f_{k+1}(R'_1, R_{-1}) = a_k$. This implies $\delta(R_1, f(R'_1, R_{-1})) - \delta(R_1, f(R_N)) = 1$.
2. $k \in P$. In this case, $f_l(R_N) = f_l(R'_1, R_{-1})$ for all $l \notin \{k, k+1\}$. In both profiles R_N and (R'_1, R_{-1}) the relative order of adjacent alternatives (a_k, a_{k+1}) is decided by committee \mathcal{C}_k . Eq. (3) and the definition of committees imply that agent 1's misreport can never be profitable.

3. $k + 1 \in P$. In this case, $f_l(R_N) = f_l(R'_1, R_{-1})$ for all $l \notin \{k, k + 1, k + 2\}$. So let us focus on those three ranks and the alternatives that occupy them. As $k \notin P$, we have $f_k(R_N) = a_k$ and $f_k(R'_1, R_{-1}) = a_{k+1}$. Subsequently, we focus on alternatives occupying ranks $k + 1$ and $k + 2$. In profile R_N , the rule f determines the relative order of adjacent alternatives (a_{k+1}, a_{k+2}) via the committee \mathcal{C}_{k+1} whereas in profile (R'_1, R_{-1}) , the rule f determines the relative order of (a_k, a_{k+2}) via the committee \mathcal{C}_{k+1} . The most profitable misreport occurs when committee \mathcal{C}_{k+1} ranks a_{k+2} before a_{k+1} (meaning $f_{k+1}(R_N) = a_{k+2}$, $f_{k+2}(R_N) = a_{k+1}$) and committee \mathcal{C}_{k+1} ranks a_k before a_{k+2} (meaning $f_{k+1}(R'_1, R_{-1}) = a_k$, $f_{k+2}(R'_1, R_{-1}) = a_{k+2}$). In that case, truthful reporting results in the triplet $a_k a_{k+2} a_{k+1}$ occupying ranks $k, k + 1, k + 2$, whereas misreporting in the triplet $a_{k+1} a_k a_{k+2}$ in those same ranks. Putting together the various possible outcomes of committees \mathcal{C}_{k+1} and \mathcal{C}_{k+1} implies $\delta(R_1, f(R'_1, R_{-1})) - \delta(R_1, f(R_N)) \in \{0, 1, 2\}$.




Figure 6: An illustration of Case 3. Orderings $f(R_N)$ and $f(R'_1, R_{-1})$ are identical at all ranks $l \notin \{k, k + 1, k + 2\}$. Committee \mathcal{C}_{k+1} determines the alternatives occupying ranks $k + 1$ and $k + 2$.

4. $k - 1 \in P$. In this case, $f_l(R_N) = f_l(R'_1, R_{-1})$ for all $l \notin \{k - 1, k, k + 1\}$. So let us focus on those three ranks and the alternatives that occupy them. As $k + 1 \notin P$, we have $f_{k+1}(R_N) = a_{k+1}$ and $f_k(R'_1, R_{-1}) = a_k$. Subsequently, we focus on alternatives occupying ranks $k - 1$ and k . In profile R_N , the rule f determines the relative order of adjacent alternatives (a_{k-1}, a_k) via the committee \mathcal{C}_{k-1} whereas in profile (R'_1, R_{-1}) , the rule f determines the relative order of (a_{k-1}, a_{k+1}) via the committee \mathcal{C}_{k-1} . The most profitable misreport occurs when committee \mathcal{C}_{k-1} ranks a_k before a_{k-1} (meaning $f_{k-1}(R_N) = a_k$, $f_k(R_N) = a_{k-1}$) and committee \mathcal{C}_{k-1} ranks a_{k-1} before a_{k+1} (meaning $f_{k-1}(R'_1, R_{-1}) = a_{k-1}$, $f_k(R'_1, R_{-1}) = a_{k+1}$). In that case, truthful reporting results in the triplet $a_k a_{k-1} a_k$ occupying ranks $k - 1, k, k + 1$, whereas misreporting in the triplet $a_{k-1} a_{k+1} a_k$ in those same ranks. Putting together the various possible outcomes of committees \mathcal{C}_{k-1} and \mathcal{C}_{k-1} implies $\delta(R_1, f(R'_1, R_{-1})) - \delta(R_1, f(R_N)) \in \{0, 1, 2\}$.




Figure 7: An illustration of Case 4. Orderings $f(R_N)$ and $f(R'_1, R_{-1})$ are identical at all ranks $l \notin \{k-1, k, k+1\}$. Committee \mathcal{C}_{k-1} determines the alternatives occupying ranks $k-1$ and k .

□

Proof of Proposition 3:

We begin with part (1). Suppose g is a Condorcet-Kemeny rule with ordering \succeq . Let $A = \{a, b, c\}$ and $R_N \in \mathcal{R}^N$. We will first argue that if K-SP is violated, then the ordering \succeq must fail regularity.

Suppose, without loss of generality, that agent i 's preferences are given by $R_i = abc$ and that there exists $R'_i \in \mathcal{R}$ such that $\delta(R_i, g(R_N)) > \delta(R_i, g(R'_i, R_{-i}))$. We distinguish four cases:

- (i) $\delta(R_i, g(R_N)) = 0$. But since $0 \leq \delta(R_i, R)$ for all $R \in \mathcal{R}$, we immediately reach a contradiction.
- (ii) $\delta(R_i, g(R_N)) = 1$. Then, we must have $\delta(R_i, g(R'_i, R_{-i})) = 0$. Hence, $R_i = g(R'_i, R_{-i})$. This implies that rule g is not Btw-SP which contradicts Proposition 5 in [17].
- (iii) $\delta(R_i, g(R_N)) = 3$. Then, we must have $\delta(R_i, g(R'_i, R_{-i})) < 3$. Let \tilde{R}_i denote the ordering which is exactly the opposite of R_i (which reverses the direction of all binary comparisons). Then, it must be the case that $g(R_N) = \tilde{R}_i$ and $g(R'_i, R_{-i}) \neq \tilde{R}_i$. This again contradicts the Btw-SP of g .
- (iv) $\delta(R_i, g(R_N)) = 2$. This is the only nontrivial case and we address it in what follows.

To violate K-SP we must have $\delta(R_i, g(R'_i, R_{-i})) < 2$. Suppose, first, that $\delta(R_i, g(R'_i, R_{-i})) = 0$. Repeating the argument of case (ii), we arrive at a contradiction.

Thus, we must have $\delta(R_i, g(R'_i, R_{-i})) = 1$. Now, $\delta(R_i, g(R_N)) = 2$ implies that we must have either $g(R_N) = cab$ or $g(R_N) = bca$. Suppose that $g(R_N) = cab$ (the proof for case

$g(R_N) = bca$ is similar). Then, to avoid violating Btw-SP we must have $g(R'_i, R_{-i}) = bac$. We will argue how this cannot happen unless the ordering \succeq violates regularity.

Given profile R_N , define the 3×3 matrix E , where E_{xy} denotes the number of agents ranking alternative x over y . For all pairs $(x, y) \in A \times A$ such that $x \neq y$ we must have $E_{xy} + E_{yx} = |N|$ (while the diagonal elements of E are set equal to 0). Hence, matrix E tabulates the results of all head-to-head contests between two alternatives under truthful preferences. Now, denote by E' the altered matrix w.r.t. to E , in which agent i misreports her true preferences $R_i = abc$ by submitting $R'_i \neq R_i$. We have the following five possibilities:

- (I) $R'_i = bac$, implying $E'_{ab} = E_{ab} - 1$, $E'_{ca} = E_{ca}$, $E'_{cb} = E_{cb}$;
- (II) $R'_i = bca$, implying $E'_{ab} = E_{ab} - 1$, $E'_{ca} = E_{ca} + 1$, $E'_{cb} = E_{cb}$;
- (III) $R'_i = acb$, implying $E'_{ab} = E_{ab}$, $E'_{ca} = E_{ca}$, $E'_{cb} = E_{cb} + 1$.
- (IV) $R'_i = cba$, implying $E'_{ab} = E_{ab} - 1$, $E'_{ca} = E_{ca} + 1$, $E'_{cb} = E_{cb} + 1$.
- (V) $R'_i = cab$, implying $E'_{ab} = E_{ab}$, $E'_{ca} = E_{ca} + 1$, $E'_{cb} = E_{cb} + 1$.

Now, since $g(R_N) = cab$ and $g(R'_i, R_{-i}) = bac$, it must be the case that:

$$E_{ca} + E_{cb} + E_{ab} \geq E_{ac} + E_{bc} + E_{ba} \quad (10)$$

$$E'_{ca} + E'_{cb} + E'_{ab} \leq E'_{ac} + E'_{bc} + E'_{ba}. \quad (11)$$

Given agent i 's five possible modifications to matrix E listed above, the only way that (10)-(11) do not lead to a contradiction is if either case (I) or (II) applies.²⁴ If case (II) applies then we must have $E_{ca} + E_{cb} + E_{ab} = E'_{ca} + E'_{cb} + E'_{ab}$ in turn implying that both (10)-(11) are equalities. But then we cannot have $g(R_N) = cab$ and $g(R'_i, R_{-i}) = bac$ (this would imply that $cab \succeq bac \succeq cab$, a contradiction).

Thus it must be that case (I) applies. Since $g(R_N) = cab$ we must have $E_{ab} \geq E_{ba}$ (otherwise, $f(R_N) \neq cab$ because ordering cba would have better Kemeny performance for profile R_N). For similar reasons, we must also have $E_{ca} + E_{cb} \geq E_{ac} + E_{bc}$, and $E_{ca} \geq E_{ac}$.

We now distinguish between two cases:

1. $E_{ca} + E_{cb} > E_{ac} + E_{bc}$. In this case we cannot have $bac \neq f(R'_i, R_{-i})$, since ordering cba would have a better Kemeny score for profile (R'_i, R_{-i}) .
2. $E_{ca} + E_{cb} = E_{ac} + E_{bc}$. Here, suppose first that $E_{ca} > E_{ac}$. Then we cannot have $bac \in K(R'_i, R_{-i})$ since bca would have better Kemeny performance for profile (R'_i, R_{-i}) .

²⁴Recall that pairs of elements symmetric to the main diagonals of E and E' must sum to $|N|$.

Hence, it must be that $E_{ac} = E_{ca}$ implying $E_{cb} = E_{bc}$. Thus, $|N|$ must be even. If $E_{ab} = E_{ba}$, then \succeq must rank cab first, and bac before cba or bca . If $E_{ab} = E_{ba} + 2$, then \succeq must rank bac first, and cab before abc or acb . In either case, the ordering \succeq is not regular.

Now, suppose that \succeq fails regularity. For ease of exposition and without loss of generality, suppose the first-ranked ordering of \succeq is cab and $bac \succeq cba$ and $bac \succeq bca$ and consider the associated \succeq -Condorcet-Kemeny rule (call it g). Construct a profile R_N such that $E_{ac} = E_{ca}$, $E_{cb} = E_{bc}$, and $E_{ab} = E_{ba}$ and where there exists an agent i with preferences $R_i = abc$. We will have $g(R_N) = cab$. Suppose this agent misreports by submitting $R'_i = bac$, leading to $g(R'_i, R_{-i}) = bac$ and implying that the rule is not K-SP.

Now we address part (2). Suppose g is a \succeq -fixed-benchmark rule. Without loss of generality, suppose that $R_i = abc$ and agent i can profitably Kemeny misreport. Since f satisfies Btw-SP by [7], the only way that K-SP can be violated is if $f(R_N) = cab$ and $f(R'_i, R_{-i}) = bac$ or $f(R_N) = bca$ and $f(R'_i, R_{-i}) = acb$. Suppose that the former case holds (the latter is handled with a similar argument).

Since f satisfies Btw-SP, we will have $f(R'_i, R_{-i}) = bac \Rightarrow f(bac, R_{-i}) = bac$. Now distinguish between the following two cases:

- (i) $bac \succeq cab$. Here, $f(R_N) = cab$ implies that bac violated strong unanimity in profile R_N . Since $(b, a) \notin R_i$, this means that either $(c, b) \in \bigcap_{i \in N} R_i$ or $(c, a) \in \bigcap_{i \in N} R_i$ or both. But this contradicts $R_i = abc$.
- (ii) $cab \succeq bac$. Here, $f(R_N) = cab$ implies that there exist $j, k \neq i$ such that $(c, a) \in R_j$ and $(c, b) \in R_k$. Since $f(bac, R_{-i}) \neq cab = f(abc, R_{-i})$, this means that $bac \cap \bigcap_{j \neq i} R_j = \{(b, a)\}$ and $abc \cap \bigcap_{j \neq i} R_j = \emptyset$. Hence, ordering cab is ranked first and $f(bac, R_{-i}) = bac$ implies that $bac \succeq bca$ and $bac \succeq cba$. Thus, \succeq is not regular.

Now, suppose we have a \succeq -fixed-benchmark rule, call it f , such that \succeq is not regular. For ease of exposition, and without loss of generality, suppose that \succeq ranks cab first and bac before both bca and cba . Consider now the profile R_N , where $R_i = abc$, $(b, a) \in R_j$ for all $j \neq i$ and $\bigcap_{l \in N} R_l = \emptyset$. Then $g(R_N) = cab$. Now, suppose agent i misreports and submits $R'_i = bac$, leading to $bac \cap \bigcap_{j \neq i} R_j = \{(b, a)\}$. The ordering \succeq ensures that $g(R'_i, R_{-i}) = bac$ leading to a violation of K-SP. \square

Proof of Theorem 7: Consider two agents and three alternatives, say $N = \{1, 2\}$ and $A = \{a, b, c\}$.

Consider the opposite profile $R_N = (R_1, R_2) = (abc, cba)$. By preference selection, $f(R_N) \in \{R_1, R_2\}$, say $f(R_N) = cba$. Now by K-SP and preference selection, $f(abc, bca) = bca$. Again by K-SP and preference selection we have $f(acb, bca) = bca$. Now applying the same arguments repetitively we obtain $f(cba, abc) = abc$ which is now a contradiction to anonymity (as $f(abc, cba) = cba$).

Now for arbitrary number m of alternatives, we may enlarge the above profiles by letting all agents rank the same $m-3$ alternatives in the same order at the bottom and by preference selection $f(R_N)$ has to rank those alternatives at the bottom with the same ranking. But then we can do the same arguments as above.

For an arbitrary number n of agents, if n is even then half of the agents play the role of agent 1 and half of the agents play the role of agent 2 and this induces a K-SP, preference selection and anonymous rule, a contradiction to the above. \square

Before proceeding with the proof of Theorem 8, we consider the special case of three alternatives and three agents and show Proposition 4.

Proof of Proposition 4: For part (i) it is easy to verify that the described rules satisfy all the properties.

For the other direction, we first show $f(abc, cba, cba) = cba$. If $f(abc, cba, cba) \neq cba$, then by preference selection, $f(abc, cba, cba) = abc$. Then by neutrality, $f(cba, abc, abc) = cba$. Now using both preference selection and K-SP it can be shown that $f(bac, cab, cab) = bac$, $f(acb, bca, bca) = acb$ and f is dictatorial with dictator 1, i.e. $f(R_N) = R_1$ for all $R_N \in \mathcal{R}^N$, a contradiction. Hence, $f(abc, cba, cba) = cba$. Similarly, we obtain $f(cba, abc, cba) = cba$ and $f(cba, cba, abc) = cba$. Using preference selection and K-SP we then obtain for any preference $R \in \mathcal{R}$ and any profile $R_N \in \mathcal{R}^N$ such that for $N = \{i, j, k\}$ we have $R_i = R_j = R$ and $R_k = -R$, $f(R_N) = R$. Furthermore, by preference selection $f(abc, cab, bca) \in \{abc, cab, bca\}$, say $f(abc, cab, bca) = abc$, and then by neutrality for all $R_N \in \Delta'$, $f(R_N) = R_1$, and similar for the triangular profiles in Δ'' . Now it is easy to see that f is a median rule with agent-based tie-breaking.

For part (ii) it is easy to verify that the described rules satisfy all the properties.

For the other direction, note that by anonymity, for any $R_N, R'_N \in \hat{\Delta}'$, $f(R_N) = f(R'_N) \equiv \hat{R}_0'$, and for any $R_N, R'_N \in \hat{\Delta}''$, $f(R_N) = f(R'_N) \equiv \hat{R}_0''$. For any $R \in \mathcal{R}^N \setminus \Delta$, if the median is not chosen, then by preference selection and anonymity, say $f(R) = R_1 \neq R_2, R_3$ and $R_1 \notin [R_2, R_3]$. Using K-SP and preference selection, then without loss of generality,

$f(abc, acb, cab) = abc$. Applying K-SP and preference selection, we obtain $f(bac, acb, cab) = bac$ and $f(bac, acb, cba) = bac$. On the other hand we may use K-SP and preference selection repetitively (by moving agent 1 first and then agents 2 and 3) to obtain $f(cab, cba, bca) = cab$. But now $f(acb, cba, bca) = acb$ and by K-SP, $f(acb, cba, bac) \neq bac$. This is a contradiction to anonymity as $f(bac, acb, cab) = bac$. \square

Proof of Theorem 8: Let $N = \{1, 2, 3\}$ and $A = \{a, b, c, d\}$. Suppose that f satisfies K-SP, preference selection and anonymity. Let \underline{f} defined for profiles where all agents rank d at the bottom. By Proposition 4 (ii), \underline{f} must be median rule with preference-based tie-breaking.

Consider the following profile $R_N = (R_1, R_2, R_3) = (abcd, cabd, bcad)$. Without loss of generality (by preference selection), let $\underline{f}(R_N) = R_1 = f(R_N)$. Consider the profile $\hat{R}_N = (\hat{R}_1, \hat{R}_2, \hat{R}_3) = (dabc, abdc, badc)$.

Starting from profile R_N we have for $R'_1 = abdc$ by K-SP and preference selection $f(R'_1, R_2, R_3) = R'_1$ (as $\delta(R'_1, R_1) = 1$, $\delta(R'_1, R_2) > 1$ and $\delta(R'_1, R_3) > 1$). Similarly, for $R''_1 = adbc$ we obtain $f(R''_1, R_2, R_3) = R''_1$ and finally $f(\hat{R}_1, R_2, R_3) = \hat{R}_1$. Then we have $\delta(R_3, \hat{R}_1) = 5 > \max\{\delta(R_3, R_2), \delta(R_3, \hat{R}_3)\}$, and from K-SP and preference selection we obtain $f(\hat{R}_1, R_2, \hat{R}_3) = \hat{R}_1$. Similarly, we then have $\delta(R_2, \hat{R}_1) = 5 > \max\{\delta(R_2, \hat{R}_3), \delta(R_2, \hat{R}_2)\}$, and from K-SP and preference selection we obtain $f(\hat{R}_1, \hat{R}_2, \hat{R}_3) = \hat{R}_1$. At profile \hat{R}_N all agents rank c at the bottom and $f(\hat{R}_N)$ restricted to $\{a, b, d\}$ shall be the median of \hat{R}_N restricted to $\{a, b, d\}$, but $f_1(\hat{R}_N) = d$ and agents 2 and 3 rank d last among $\{a, b, d\}$, a contradiction (as the median rule would require $f_3(\hat{R}_N) = d$).

Now if $|N| = 3k$ and $m \geq 4$, then k agents play the role of agent 1 (by reporting the same preference), k agents play the role of agent 2 and k agents play the role of agent 3 and we obtain a three-agents rule satisfying K-SP, preference selection and AN. By letting all agents rank the same $m - 4$ alternatives at the bottom, then we obtain a contradiction as above (since by preference selection those alternatives must be ranked at the bottom of the social ranking as well). \square

References

- [1] Alemante, F., Campbell, D. E., and Kelly, J. S. (2016). Characterizing the resolute part of monotonic social choice correspondences. *Economic Theory*, 62, 765-783.
- [2] Amorós, P. (2009). Eliciting optimal rankings from unfair jurors. *Journal of Economic Theory*, 144, 1211-1226.
- [3] Amorós, P., Corchón, L. C., and Moreno, B. (2002). The scholarship assignment problem. *Games and Economic Behavior*, 38, 1-18.

- [4] Arrow, K. J. (1959). Rational choice functions and orderings. *Economica*, 26, 121-127.
- [5] Arrow, K.J. (1963). *Social choice and individual values*. 2nd edition. John Wiley. New York.
- [6] Athanassoglou, S. (2016). Strategyproof and efficient preference aggregation with Kemeny-based criteria. *Games and Economic Behavior*, 95, 156-167.
- [7] Athanassoglou, S. (2019). Solidarity and efficiency in preference aggregation: a tale of two rules, *Games and Economic Behavior*, 118, 126-140.
- [8] Aziz, H., Lam, A., Lee, B. E., and Walsh, T. (2021). Strategyproof and proportionally fair facility location. arXiv preprint arXiv:2111.01566.
- [9] Barberà, S. (2011). Strategyproof social choice. In: Arrow, K.J., Sen, A., Sen K. (Eds), *Handbook of Social Choice and Welfare*, vol 2. Elsevier, Amsterdam, pp. 731-831. Chapter 25.
- [10] Barberà, S., Berga, D., and Moreno, B. (2012). Group strategy-proof social choice functions with binary ranges and arbitrary domains: characterization results. *International Journal of Game Theory*, 41, 791-808.
- [11] Barberà, S., Berga, D., Moreno, B., and Nicolò, A. (2023). Condorcet consistency and pairwise justifiability under variable agendas. Working Paper.
- [12] Barberà, S., Gul, F., and Stacchetti, E. (1993). Generalized median voter schemes and committees. *Journal of Economic Theory*, 61(2), 262-289.
- [13] Barberà, S., Massó, J., and Neme, A. (1997). Voting under constraints. *Journal of Economic Theory*, 76(2), 298-321.
- [14] Barberà, S., Sonnenschein, H., and Zhou, L. (1991). Voting by committees. *Econometrica*, 59, 595-609.
- [15] Black, D. (1948). On the rationale of group decision making. *Journal of Political Economy*, 56, 23-34.
- [16] Bonkougou, S. (2018). Strategy-proof preference aggregation: augmented serial rules. Working Paper.
- [17] Bossert, W., and Sprumont, Y. (2014). Strategy-proof preference aggregation: possibilities and characterizations. *Games and Economic Behavior*, 85, 109-126.
- [18] Bartholdi, L., Hann-Caruthers, W., Josyula, M., Tamuz, O., and Yariv, L. (2021). Equitable voting rules. *Econometrica*, 89(2), 563-589.
- [19] Bossert, W., and Storcken, T. (1992). Strategy-proofness of social welfare functions: the use of the Kemeny distance between preference orderings. *Social Choice and Welfare*, 9(4), 345-360.
- [20] Can, B. (2014). Weighted distances between preferences. *Journal of Mathematical Economics*, 51, 109-115.
- [21] Can, B., and Storcken, T. (2018). A re-characterization of the Kemeny distance. *Journal of Mathematical Economics*, 79, 112-116.
- [22] Dasgupta, P., and Maskin, E. (2008). On the robustness of majority rule. *Journal of the European Economic Association*, 6(5), 949-973.
- [23] Demange, G. (1982). Single-peaked orders on a tree. *Mathematical Social Sciences*, 3(4), 389-396.
- [24] Gibbard, A. (1973). Manipulation of voting schemes: a general result. *Econometrica*, 41(4), 587-601.

- [25] Ehlers, L., and Storcken, T. (2008). Arrow’s possibility theorem for one-dimensional single-peaked preferences. *Games and Economic Behavior*, 64, 533-547.
- [26] Goodin, R. E., and List, C. (2006). A conditional defense of plurality rule: generalizing May’s theorem in a restricted informational environment. *American Journal of Political Science*, 50(4), 940-949.
- [27] Harless, P. (2016). Solidarity in preference aggregation: Improving on a status quo. *Games and Economic Behavior*, 95, 73-87.
- [28] Hassanzadeh, F. F., and Milenkovic, O. (2014). An axiomatic approach to constructing distances for rank comparison and aggregation. *IEEE Transactions on Information Theory*, 60(10), 6417-6439.
- [29] Horan, S., Osborne, M. J., and Sanver, M. R. (2019). Positively responsive collective choice rules and majority rule: a generalization of May’s theorem to many alternatives. *International Economic Review*, 60(4), 1489-1504.
- [30] Kemeny, J.G. (1959). Mathematics without Numbers. *Daedalus*, 88, 575–591.
- [31] Kemeny, J.G., and Snell, J.L. (1962). Mathematical models in the social sciences (Vol. 9). New York: Blaisdell.
- [32] Kivinen, S. (2024). Equitable, neutral, and efficient voting rules. *Journal of Mathematical Economics*, 115, 103061.
- [33] Kumar, U., Roy, S., Sen, A., Yadav, S., & Zeng, H. (2021). Local-global equivalence in voting models: A characterization and applications. *Theoretical Economics*, 16(4), 1195-1220.
- [34] Kumar, U., Roy, S., Sen, A., Yadav, S., & Zeng, H. (2021). Local global equivalence for unanimous social choice functions. *Games and Economic Behavior*, 130, 299-308.
- [35] Laffond, G., Lainé, J., and Sanver, M. R. (2020). Metrizable preferences over preferences. *Social Choice and Welfare*, 55(1), 177-191.
- [36] Manjunath, V. (2012). Group strategy-proofness and voting between two alternatives, *Mathematical Social Sciences*, 63, 239-242.
- [37] May, K. O. (1952). A set of independent necessary and sufficient conditions for simple majority decision. *Econometrica*, 680-684.
- [38] Meskanen, T., & Nurmi, H. (2006). Distance from consensus: A theme and variations. In *Mathematics and Democracy: recent advances in voting systems and collective choice* (pp. 117-132). Springer Berlin Heidelberg.
- [39] Moulin, H. (1980). On strategy-proofness and single peakedness. *Public Choice*, 35(4), 437-455.
- [40] Moulin, H. (1983). *The strategy of social choice*. North-Holland.
- [41] Nishimura, H., and Ok, E. A. (2023). A class of dissimilarity semimetrics for preference relations. *Mathematics of Operations Research*, 49(4), 2249-2270.
- [42] Sato, S. (2013). A sufficient condition for the equivalence of strategy-proofness and nonmanipulability by preferences adjacent to the sincere one. *Journal of Economic Theory*, 148(1), 259-278.
- [43] Satterthwaite, M. A. (1975). Strategy-proofness and Arrow’s conditions: Existence and correspondence theorems for voting procedures and social welfare functions. *Journal of Economic Theory*, 10(2), 187-217.

- [44] Schummer, J., and Vohra, R. V. (2002). Strategy-proof location on a network. *Journal of Economic Theory*, 104(2), 405-428.
- [45] Young, H. P., and Levenglick, A. (1978). A consistent extension of Condorcet's election principle. *SIAM Journal on Applied Mathematics*, 35(2), 285-300.

Online Appendix for “Strategy-proof preference aggregation and the anonymity-neutrality tradeoff”

A Example of a non-dictatorial rule that is neutral and Min-SP but not Btw-efficient

Let $|A| \geq 3$ and $|N| \geq 2$. Given $R \in \mathcal{R}$, let $\text{bot}(R)$ (resp. $\text{top}(R)$) denote the last-ranked (resp. first-ranked) alternative of ordering R . In addition, let $R|A \setminus \{x\}$ denote the ordering R restricted to alternatives in $A \setminus \{x\}$. For $R_N \in \mathcal{R}^N$, the rule f is defined as follows:

$$f(R_N) = \begin{cases} R_1 & \text{if } R_1 = R_2, \\ R_1 & \text{if } R_1 = -R_2, \\ xR_1|A \setminus \{x\} & \text{if } \text{bot}(R_1) = \text{bot}(R_2) = x \text{ and } R_1 \neq R_2, \\ R_2 & \text{otherwise.} \end{cases}$$

It is obvious that f satisfies neutrality and violates betweenness efficiency as, for any profile such that $\text{bot}(R_i) = x$ for all $i \in N$ and $R_1 \neq R_2$, alternative x (which is the common worst alternative) is ranked at the top of $f(R_N)$. It remains to show that f satisfies Min-SP. All agents $i \in N \setminus \{1, 2\}$ have no effect on the rule's outcome. If $R_1 = R_2$, then there is nothing to show. Thus, let $R_1 \neq R_2$. We will refer to the definition of the rule as cases one to four.

First we consider agent 1. Again if $R_1 = -R_2$, then there is nothing to show. If $\text{bot}(R_1) = \text{bot}(R_2) = x$ and $R_1 \neq R_2$, then any deviation $R'_1 \neq R_1$ will lead either to the first, second, or fourth case. None of those deviations can result in $f(R'_1, R_{-1}) = R_1$.

Second, we consider agent 2. If $f(R_N)$ is determined by the fourth case (“otherwise”) there is nothing to show. If $R_1 = -R_2$, then any deviation $R'_2 \neq R_2$ will lead either to the first, third, or fourth case. None of those deviations can result in $f(R'_2, R_{-2}) = R_2$. If $\text{bot}(R_1) = \text{bot}(R_2) = x$ and $R_1 \neq R_2$, then any deviation $R'_2 \neq R_2$ will lead either to the first, second, or fourth case. None of those deviations can result in $f(R'_2, R_{-2}) = R_2$.

B Example of a rule that is anonymous, strongly unanimous and Min-SP but not Btw-SP

Let $|N| \geq 2$. Let $R \in \mathcal{R}$ denote a status-quo ordering. Consider the following rule f defined by the following iterative procedure:

- (i) Consider the alternatives ranked first by all agents and select the one ranked highest by the status quo ordering R . This alternative is put in first place by the rule and then discarded. Agent and status-quo orderings are updated accordingly.
- (ii) For every $k = 2, \dots, m$, repeat the process described in (i) to determine the k 'th-ranked alternative.

Here is an example of how this rule works. Suppose $R_1 = dbca, R_2 = cadb, R_3 = dabc$ and $R = abcd$. The highest-ranked alternatives of agents 1, 2 and 3, are d, c, d respectively. Since R ranks c higher than

d , the rule sets $f_1(R_N) = c$,²⁵ and c is removed from agent orderings and R . These are updated to $R_1^1 = dba$, $R_2^1 = adb$, $R_3^1 = dab$ and $R^1 = abd$, respectively. Now, the highest-ranked alternatives of agents 1, 2 and 3, are d, a, d respectively. So the rule sets $f_2(R_N) = a$ and a is removed from R_1^1, R_2^1, R_3^1 and R^1 . These are updated to $R_1^2 = db$, $R_2^2 = db$, $R_3^2 = db$ and $R^2 = bd$, respectively. Now since the remaining agents are unanimous in their ordering of d over b it is easy to see that the rule sets $f_3(R_N) = d$, $f_4(R_N) = b$. So, $f(R_N) = cadb$.

The rule f is clearly anonymous. To show why it is strongly unanimous, suppose $aR_i b$ for all $i \in N$. If f ranked b higher than a , then there would have to be an agent whose current highest-ranked ordering is b . But this is a contradiction.

We now show that f is Min-SP but not Btw-SP. Suppose that $f(R_N) \neq R_i$, and let $k \in \{1, \dots, m\}$ be the first rank at which R_i and $f(R_N)$ differ. Suppose agent i 's k 'th-ranked alternative is b_k . Hence, there exists $b \neq b_k$ and $j \neq i \in N$ such that b is ranked higher than b_k by R and b is agent j 's top-ranked alternative among those left. We claim that there exists no R'_i such that $f_l(R'_i, R_{-i}) = f_l(R_N)$ for all $l < k$, and $f_k(R'_i, R_{-i}) = b_k$. Consider a preference deviation $R'_i \neq R_i$ that maintains the identity and order of the first $k-1$ alternatives. In order for $f_k(R'_i, R_{-i}) \neq f_k(R_N)$, agent i has to misreport his current highest-ranked alternative and the rule f must pick it. Hence, $f_k(R'_i, R_{-i}) \neq b_k$ and $f(R'_i, R_{-i}) \neq f(R_N)$. We conclude that the f is Min-SP.

To see why f is not Btw-SP, consider $R_1 = dbca$, $R_2 = abcd$ and $R = bacd$. We have $f(R_N) = abcd$. Let $R'_1 = bdac$. Then $f(R'_1, R_2) = bacd \in [dbca, abcd]$.

C Example of a rule that is anonymous, neutral, strongly unanimous and Min-SP but not Btw-SP

Let $A = \{a, b, c\}$ and $N = \{1, \dots, 7\}$. For any $R \in \mathcal{R}^N$, $f(R)$ is defined as follows: the top ranked alternative of $f(R)$ is the alternative which is top ranked by most agents, where if there is a tie, it must be that three agents rank x at the top and three agents rank y at the top, then the unique agent who ranks neither x nor y at the top determines whether x or y is ranked at the top of $f(R)$; and afterwards the remaining two alternatives are ranked according to majority. Note that f is well-defined as tie for the top ranked alternatives (as there are three alternatives and seven agents) only if three agents rank x at the top, three agents rank y at the top, and the remaining agent ranks z at the top (where $A = \{x, y, z\}$). After having determined the top ranked alternative of $f(R)$, majority relation determines the ranking of the remaining two alternatives as the number of agents is odd.

It is obvious that f satisfies anonymity, neutrality and strong unanimity. We show that f satisfies Min-SP. Consider agent i and a deviation $R' = (R'_i, R_{-i})$ from R . Now if $\text{top}(f(R)) = \text{top}(f(R'))$, then the deviation is not profitable for agent i as majority over two remaining alternatives is non-manipulable. Thus, let $\text{top}(f(R)) \neq \text{top}(f(R'))$. But then if Min-SP is violated, we must have $\text{top}(f(R')) = \text{top}(R_i) \neq \text{top}(f(R))$. If four agents or more (other than i) have $\text{top}(f(R))$ as the top ranked alternative, then agent i cannot change the top ranked alternative at $f(R')$, and also this manipulation is impossible if agent i is the unique agent with a different top ranked alternative (as then he cannot change the socially top ranked alternative to his favor). For the remaining case, three agents rank x at the top, two agents rank y at the top, and two agents

²⁵As in the paper, $f_l(R_N)$ denotes the l -ranked alternative of ordering $f(R_N)$.

rank z at the top. But then agent i ranks either y or z at the top and it is impossible for him to change the socially top ranked alternative in his favor.

In order to see that f violates betweenness-SP, let $R_1 = R_2 : abc$, $R_3 = R_4 : bac$ and $R_5 = R_6 = R_7 : cab$. Then $f(R) : cab$. Now let $R'_3 : abc$ and $R' = (R'_3, R_{-3})$. Then $f(R') : abc$, $f(R') \in [R_3, f(R)]$ and $f(R') \neq f(R)$, i.e. f violates betweenness-SP.

Another undesirable feature of the rule f is that it might a Condorcet loser at the top of its ranking: for $R_1 = R_2 : bac$, $R_3 = R_4 : abc$ and $R_5 = R_6 = R_7 : cba$ we have $f(R) : cba$ while four agents prefer both a and b over c , i.e. c is a Condorcet loser.

D Example of a rule that is Local K-SP but not Min-SP

Suppose $m = 4$ and consider a function $f : \mathcal{R} \mapsto \mathcal{R}$ satisfying:

$$\begin{aligned} f(abcd) &= adbc, & f(abdc) &= badc, & f(acbd) &= acdb, & f(adbc) &= adcb \\ f(bacd) &= badc, & f(bdca) &= bcda, & f(cbad) &= cabd, & f(cdab) &= dcab \\ f(dbca) &= dabc, & f(dcba) &= cdab, & f(dacb) &= dcab, & f(dbac) &= dabc \\ f(R) &= R, \text{ for all other } R \in \mathcal{R} \end{aligned}$$

Define the rule $g : \mathcal{R}^N \mapsto \mathcal{R}$ such that for all $R_N \in \mathcal{R}^N$, we have $g(R_N) = f(R_1)$. To verify its stated properties we only need to check deviations by agent 1. To check Local K-SP it suffices to examine the profitability of adjacent deviations for agent 1 when R_1 is such that $f(R_1) \neq R_1$. There are $12 \cdot 3 = 36$ such deviations and none of them imply a violation of Local K-SP. For example, when $R_1 = abcd$, we have $g(R_N) = f(abcd) = adbc$, $\delta(R_1, g(R_N)) = 2$, and three possible local misreports: $R'_1 \in \{bacd, acbd, abdc\}$. If $R'_1 = bacd$ then $g(R'_1, R_{-1}) = badc$; if $R'_1 = acbd$ then $g(R'_1, R_{-1}) = acdb$; and if $R'_1 = abdc$ then $g(R'_1, R_{-1}) = badc$. In all three cases, the Kemeny distance between R_1 and $g(R'_1, R_{-1})$ is exactly equal to 2.

On the other hand, if $R_1 = adbc$, and agent 1 misreports $R'_1 = abcd$, then $g(R_N) \neq R_1$ and $g(R'_1, R_{-1}) = R_1$, in violation of Min-SP. Note that $\delta(R_1, R'_1) = 2$, so that this violation of Min-SP has no bearing on Local K-SP. Note, finally, that the rule g is not onto (e.g., there is no profile R_N such that $g(R_N) = abcd$).

E Integer programming formulation and application to $n = 2, m = 4$ case

A rule f is a function $f : \mathcal{R}^N \mapsto \mathcal{R}$, determining for every profile a social ordering. If a rule f , when applied to profile R_N , ranks a_i before a_j then we write $a_i f(R_N) a_j$. Equivalently, we may introduce the binary variables $d_{R_N}^f(a_i, a_j) \in \{0, 1\}$ and require that :

$$d_{R_N}^f(a_i, a_j) = 1 \Leftrightarrow a_i f(R_N) a_j.$$

In this way we can express f by associating with it a vector $\mathbf{d}^f \equiv [d_{R_N}^f(x, y)]$ where R_N ranges throughout \mathcal{R}^N and $(a_i, a_j) \in A \times A$ with $i \neq j$. As a result \mathbf{d}^f is a vector of dimension $(m!)^n \cdot m(m-1)$ that must satisfy the binary integer requirement $d_{R_N}^f(a_i, a_j) \in \{0, 1\}$ for all profiles R_N and pairs of alternatives (a_i, a_j) .²⁶

Each rule f is uniquely defined by its corresponding \mathbf{d} vector. For instance, take a dictatorial rule which, no matter what other agents submit, always selects agent 1's ordering. This rule is described by a \mathbf{d} vector such that for all profiles R_N and pairs (a_i, a_j) we have $d_{R_N}(a_i, a_j) = 1$ if and only if $(a_i, a_j) \in R_1$.

We define the rule properties that were discussed in the main text, each time translating them to the integer-vector framework mentioned above.²⁷

- (C) **Completeness.** For all profiles $R_N \in \mathcal{R}^N$ and pairs of alternatives $(a_i, a_j) \in A \times A$ and $i \neq j$, either $a_i f(R_N) a_j$ or $a_j f(R_N) a_i$.

Equivalently, for all $R_N \in \mathcal{R}^N$:

$$d_{R_N}(a_i, a_j) + d_{R_N}(a_j, a_i) = 1, \quad \forall (a_i, a_j) \in A \times A, i \neq j. \quad (12)$$

- (T) **Transitivity.** For all profiles $R_N \in \mathcal{R}^N$ and triplets of alternatives $(a_i, a_j, a_k) \in A \times A \times A$, if $a_i f(R_N) a_j$ and $a_j f(R_N) a_k$, then $a_i f(R_N) a_k$.

Equivalently for all $R_N \in \mathcal{R}^N$:

$$d_{R_N}(a_i, a_j) + d_{R_N}(a_j, a_k) + d_{R_N}(a_k, a_i) \leq 2, \quad \forall (a_i, a_j, a_k) \in A \times A \times A, i \neq j, j \neq k, i \neq k. \quad (13)$$

- (A) **Anonymity.** Consider the set of permutations on N , and call it Σ_N . For any element $\sigma \in \Sigma_N$ and profile R_N , let $\sigma(R_N)$ denote the profile in which agent 1 is given the ordering of agent $\sigma(1)$, agent 2 of $\sigma(2)$ and so on. I.e., $\sigma(R_N) = (R_{\sigma(1)}, R_{\sigma(2)}, \dots, R_{\sigma(n)})$.

Anonymity requires that for all $R_N \in \mathcal{R}^N$ we have:

$$d_{R_N}(a_i, a_j) = d_{\sigma(R_N)}(a_i, a_j), \quad \forall \sigma \in \Sigma_N, (a_i, a_j) \in A \times A, i \neq j. \quad (14)$$

- (N) **Neutrality.** Consider now the set of permutations on A , call it Π_A . For any element $\pi \in \Pi_A$ and profile R_N , let $\pi(R_N)$ denote the profile in which agents' rankings are rewritten in a way that alternative a_1 is now named $\pi(a_1)$, alternative a_2 is named $\pi(a_2)$ and so on. That is, $\pi(R_N) = (\pi(R_1), \pi(R_2), \dots, \pi(R_n))$, where $\pi(R_k) = \{(\pi(a_i), \pi(a_j)) : \forall (a_i, a_j) \in R_k\}$. Neutrality requires that for all $R_N \in \mathcal{R}^N$ we have:

$$d_{R_N}(a_i, a_j) = d_{\pi(R_N)}(\pi(a_i), \pi(a_j)), \quad \forall \pi \in \Pi_A, (a_i, a_j) \in A \times A, i \neq j. \quad (15)$$

- (U) **Unanimity.** If all agents submit the same ordering then the rule also assigns this ordering.

Equivalently, for all $R_N \in \mathcal{R}_N$ and $R \in \mathcal{R}$:

$$\{R_k = R, \forall k = 1, 2, \dots, n\} \Rightarrow \{d_{R_N}(a_i, a_j) = 1, \forall (a_i, a_j) \in R\}. \quad (16)$$

²⁶In what follows for convenience we drop the superscript f from the vector \mathbf{d}^f .

²⁷Completeness and Transitivity are explicitly listed because the integer-vector formulation does not a priori ensure they hold.

(SU) **Strong Unanimity.** If all agents prefer an alternative a_i to a_j then the rule must also respect this binary comparison.

Equivalently, for all $R_N \in \mathcal{R}^N$ and $(a_i, a_j) \in A \times A$:

$$\{a_i R_k a_j, \forall k = 1, 2, \dots, n\} \Rightarrow d_{R_N}(a_i, a_j) = 1. \quad (17)$$

Clearly, LU implies the U property above.

(K-SP) **Kemeny strategy-proofness.** Consider a profile $R_N = (R_1, R_2, \dots, R_n)$ and within this profile an agent k with preferences R_k . Denote by R_{-k} the preferences of all other agents, i.e., $R_{-k} = (R_1, R_2, \dots, R_{k-1}, R_{k+1}, \dots, R_n)$. Suppose now agent k misreports his preferences by submitting an ordering $R'_k \neq R_k$. This will give rise to a profile $(R'_k, R_{-k}) \equiv (R_1, R_2, \dots, R_{k-1}, R'_k, R_{k+1}, \dots, R_n)$. K-SP requires that for all $R_N \in \mathcal{R}^N$ and $k \in N$ all such misreports not be profitable so that:

$$\sum_{(a_i, a_j) \in R_k} d_{R_N}(a_i, a_j) \geq \sum_{(a_i, a_j) \in R_k} d_{(R'_k, R_{-k})}(a_i, a_j), \quad \forall R'_k \neq R_k. \quad (18)$$

The Integer Program. Set $A = \{a, b, c, d\}$ and $N = \{1, 2\}$. We wish to explore whether there exists a rule satisfying Anonymity, Strong Unanimity and K-SP.

Using an integer-programming framework, this problem can be reformulated in the following way: Does there exist a 0-1 \mathbf{d} vector that simultaneously satisfies (C)-(T)-(A)-(SU)-(K-SP)? Since $m = 4, n = 2$, the associated \mathbf{d} vector will have dimension $(4!)^2 * 4 * 3 = 6912$. To answer this question, we set up an integer program. One particularly simple way of doing this is to we introduce a scalar variable $y \geq 0$ and define problem \mathcal{P} :

$$\begin{aligned} \mathcal{P} = \min_{\mathbf{d}, y} \quad & y \\ \text{s.t.} \quad & \text{(C)-(T)-(A)-(SU)} \\ & \sum_{(a_i, a_j) \in R_k} d_{R_N}(a_i, a_j) - \sum_{(a_i, a_j) \in R_k} d_{(R'_k, R_{-k})}(a_i, a_j) + y \geq 0, \quad \forall R_N \in \mathcal{R}^N, k \in N, R'_k \neq R_k. \\ & \mathbf{d} \in \{0, 1\}, y \geq 0. \end{aligned}$$

If \mathcal{P} yields an optimal solution (\mathbf{d}^*, y^*) with $y^* > 0$, we will know that anonymity, strong unanimity and K-SP cannot be simultaneously satisfied. Conversely, if $y^* = 0$ then the vector \mathbf{d}^* defines a rule that satisfies the desired properties. It is thus clear that finding an optimal solution for \mathcal{P} is equivalent to resolving the existence of an anonymous, strongly unanimous, K-SP rule.

So our goal is to solve mixed-integer program \mathcal{P} or closely-related variants thereof. To this end, we use Matlab and its built-in mixed-integer linear programming solver **intlinprog** to solve a more involved version of \mathcal{P} in which there is a non-negative variable y_h for each K-SP constraint h ,²⁸ and a linear objective function $\mathbf{c} \cdot \mathbf{y}$, where $\mathbf{c} > \mathbf{0}$. We do this for computational reasons, as the simpler model seemed to be yielding suboptimal results.

²⁸There are $(4!)^2 * 2 * 23 = 26496$ such constraints.

$$\begin{aligned} \mathcal{P}' = \min_{\mathbf{d}, \mathbf{y}} \quad & \mathbf{c} \cdot \mathbf{y} \\ \text{s.t.} \quad & \text{(C)-(T)-(A)-(SU)} \end{aligned}$$

$$\begin{aligned} \sum_{(a_i, a_j) \in R_k} d_{R_N}(a_i, a_j) - \sum_{(a_i, a_j) \in R_k} d_{(R'_k, R_{-k})}(a_i, a_j) + y_h \geq 0, \quad \forall R_N \in \mathcal{R}^N, k \in N, R'_k \neq R_k. \\ \mathbf{d} \in \{0, 1\}, \quad \mathbf{y} \geq \mathbf{0}. \end{aligned}$$

Problem \mathcal{P}' shares the appealing properties of \mathcal{P} : the existence of an anonymous, strongly unanimous, K-SP rule is equivalent to an optimal solution $(\mathbf{d}^*, \mathbf{y}^*)$ of \mathcal{P}' such that $\mathbf{y}^* = \mathbf{0}$. After a few iterations and various modifications of vector \mathbf{c} , the solver converges to the rule discussed in the text and listed in the Excel file, and we have a proof of existence.

Below we give the complete formal definition of the rule found as a solution to the above problem. First, we define fixed-status-quo rules with tie-breaking for three alternatives.

Definition 11 Let $N = \{1, 2\}$ and $A = \{a_1, a_2, a_3\}$. Fix a status-quo ordering R_0 and denote by $r_k(R)$ the k 'th-ranked alternative of $R \in \mathcal{R}$. The R_0 -fixed-status-quo rule with tie-breaking, SQ^{R_0} , is the anonymous rule such that for all $R_N = (R_1, R_2) \in \mathcal{R}^N$:

- (i) if R_0 is strongly unanimous for R_N , then $SQ^{R_0}(R_N) = R_0$;
- (ii) if R_0 is not strongly unanimous for R_N and $R_1 \in [R_0, R_2]$, then $SQ^{R_0}(R_N) = R_1$;
- (iii) if R_0 is not strongly unanimous for R_N and $R_2 \in [R_0, R_1]$, then $SQ^{R_0}(R_N) = R_2$; and
- (iv) otherwise, we have $R_1 \neq R_2$ and $\delta(R_1, R_0) = \delta(R_2, R_0) = 2$, and we set $SQ^{R_0}(R_N) = R_i$, where $i \in N$ is such that $r_3(R_i) = r_1(R_0)$.²⁹

It is obvious that the above rules satisfy strong unanimity and anonymity, and one can also check K-SP.

Second, for four alternatives we provide below a formal description of the anonymous, strongly unanimous and K-SP rule.

Definition 12 Let $N = \{1, 2\}$ and $A = \{a, b, c, d\}$. Specify $a \in A$ as the losing alternative. For all non-losing alternatives $x \in \{b, c, d\}$, introduce the orderings \underline{R}_0^x and \overline{R}_0^x as follows:

$$\begin{aligned} \underline{R}_0^b &= dca, & \overline{R}_0^b &= cda \\ \underline{R}_0^c &= bda, & \overline{R}_0^c &= dba \\ \underline{R}_0^d &= cba, & \overline{R}_0^d &= bca. \end{aligned}$$

Let f be an anonymous rule such that, for any profile $R_N = (R_1, R_2) \in \mathcal{R}^N$,

- (I) if $R_1 \cap R_2 = \{(b, a), (c, a), (d, a)\}$, then there are three kinds of possible profiles³⁰ and the rule sets:

$$f(bcda, dbca) = cdab, \quad f(bdca, cdab) = dbca, \quad f(cbda, dbca) = bcda;$$

²⁹To illustrate part (iv) of the definition, if $R_0 = bca$, then the R_0 -fixed-status-quo rule with tie-breaking is given by $SQ^{R_0}(abc, cab) = cab$.

³⁰Recall that the rule is anonymous.

(II) if $R_1 \cap R_2 = \{(a, b), (a, c), (a, d)\}$, then there are three kinds of possible profiles and the rule sets:

$$f(abcd, adcb) = abdc, \quad f(acbd, adbc) = adbc, \quad f(acdb, abdc) = acbd;$$

(III) if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \{(a, x)\}$ (where $x \in \{b, c, d\}$), then

$$f(R_1, R_2) = SQ^{\underline{R}_0^x}(R_{1|A \setminus \{x\}}, R_{2|A \setminus \{x\}}) x,$$

i.e., the non-losing alternative x is placed at the bottom of the social ordering and is preceded by the outcome of the \underline{R}_0^x -fixed-order-status-quo rule applied to profile $(R_{1|A \setminus \{x\}}, R_{2|A \setminus \{x\}})$;

(IV) if $R_1 \cap R_2 \cap \{(a, b), (a, c), (a, d)\} = \{(a, y), (a, z)\}$ (where $\{b, c, d\} = \{x, y, z\}$), then

$$f(R_1, R_2) = x SQ^{\overline{R}_0^x}(R_{1|A \setminus \{x\}}, R_{2|A \setminus \{x\}}),$$

i.e., the non-losing alternative x is placed at the top of the social ordering and is succeeded by the outcome of the \overline{R}_0^x -fixed-status-quo rule applied to profile $(R_{1|A \setminus \{x\}}, R_{2|A \setminus \{x\}})$;

(V) if $R_1 \cap R_2 = \emptyset$, then there are twelve possible profiles and the rule sets:

$$\begin{aligned} f(abcd, dcba) &= cdab, & f(abdc, cdab) &= dbca, & f(acbd, dbca) &= bcda, & f(acdb, bdca) &= dbca \\ f(adbc, cbda) &= bcda, & f(adcb, bcda) &= cdab, & f(bacd, dcab) &= cdab, & f(badc, cdab) &= dbca \\ f(bcda, dacb) &= cdab, & f(bdac, cadb) &= dbca, & f(cabd, dbac) &= bcda, & f(cbad, dabc) &= bcda. \end{aligned}$$

The rule's output for any other profile is given in the Excel-file but can also be derived from the fact that losing alternative a is always placed as low as possible subject to strong unanimity, from points (I)-(II)-(III)-(IV)-(V), and from properties anonymity, strong unanimity and K-SP of the rule.